

Power-Enhanced Multiple Decision Functions Controlling Family-Wise Error and False Discovery Rates¹

Edsel A. Peña²

Joshua D. Habiger³

Wensong Wu⁴

Department of Statistics
University of South Carolina
Columbia, SC 29208 USA

June 29, 2010

¹The authors acknowledge support from National Science Foundation (NSF) Grant DMS 0805809, National Institutes of Health (NIH) Grant RR17698, and Environmental Protection Agency (EPA) Grant RD-83241902-0 to University of Arizona with subaward number Y481344 to the University of South Carolina.

²E. Peña is Professor, Department of Statistics, University of South Carolina, Columbia. E-Mail: pena@stat.sc.edu.

³J. Habiger is a PhD student in the Department of Statistics, University of South Carolina, Columbia. E-Mail: habigerj@mailbox.sc.edu.

⁴W. Wu is a PhD student in the Department of Statistics, University of South Carolina, Columbia. E-Mail: wu26@mailbox.sc.edu.

Abstract

Improved procedures, in terms of smaller missed discovery rates (MDR), for performing multiple hypotheses testing with weak and strong control of the family-wise error rate (FWER) or the false discovery rate (FDR) are developed and studied. The improvement over existing procedures such as the Šidák procedure for FWER control and the Benjamini-Hochberg (BH) procedure for FDR control is achieved by exploiting possible differences in the powers of the individual tests. Results signal the need to take into account the powers of the individual tests and to have multiple hypotheses decision functions which are not limited to simply using the individual p -values, as is the case for example with the Šidák, Bonferroni, or BH procedures. They also enhance understanding of the role of the powers of individual tests, or more precisely the receiver operating characteristic (ROC) functions of decision processes, in the search for better multiple hypotheses testing procedures. A decision-theoretic framework is utilized, and through auxiliary randomizers the procedures could be used with discrete or mixed-type data or with rank-based nonparametric tests. This is in contrast to existing p -value based procedures whose theoretical validity is contingent on each of these p -value statistics being stochastically equal or greater than a standard uniform variable under the null hypothesis. Proposed procedures are relevant in the analysis of high-dimensional “large M , small n ” data sets arising in the natural, physical, medical, economic, and social sciences, whose generation and creation is accelerated by advances in high-throughput technology, notably, but not limited to, microarray technology.

Keywords and Phrases: Benjamini-Hochberg procedure; Bonferroni procedure; decision process; false discovery rate (FDR); family wise error rate (FWER) Lagrangian optimization; Neyman-Pearson most powerful test; microarray analysis; reverse martingale; missed discovery rate (MDR); multiple decision function and process; multiple hypotheses testing; optional sampling theorem; power function; randomized p -values; generalized multiple decision p -values; ROC function; Šidák procedure.

1 Introduction and Motivation

Advances in modern technology, spearheaded by microarray technology, have led to the creation or generation of many data sets characterized by a large number, M , of sets of variables, with the m th set $\mathcal{S}_m$ composed of variables which pertain to characteristics of the m th attribute of an observational unit. For historical reasons an attribute will be referred to as a ‘gene’. The variables $\mathbf{Z}_m$ in $\mathcal{S}_m$ are only measured or observed for a small number of units. Such variables may come in varied types such as being continuous, categorical, discrete, mixed, or even as functional data. They may also possess an inherent data structure such as being a multi-group data, a regression-type data, or as an event-time data with covariates and with right-censoring or truncation. Such data sets will symbolically be represented by the collection of random elements

$$\text{DATA} \equiv \{\mathbf{Z}_{mj} : j = 1, 2, \dots, n; m = 1, 2, \dots, M\}$$

with n denoting the number of units observed, i.e., the number of replications. It is typical to have $n \ll M$. To simplify notation and introduce conciseness and generality, the observables for gene m will be denoted by $X_m \equiv \{\mathbf{Z}_{mj} : j = 1, 2, \dots, n_m\}$.

Efron [12], for example, described four such data sets. The first is a prostate data set from [44] with $M = 6033$ genes and for each gene is associated a variable Z_{m1} indicating presence (value = 1) or absence (value = 0) of prostate cancer and a variable Z_{m2} representing a continuous response. For the m th gene the random vector $\mathbf{Z}_m = (Z_{m1}, Z_{m2})$ was observed on $n = 102$ replications, and these 102 subjects were utilized to compare the diseased ($Z_{m1} = 1$) and the non-diseased ($Z_{m1} = 0$) groups with respect to the response variable Z_{m2} using a two-sample t -test. The other three data sets described in [12] were an education data set from [33] with $M = 3748$; a proteomics data set from [54] with $M = 230$ and $n = 551$; and an imaging data from [40] with $M = 15445$ and $n = 12$. In all of these data sets, a decision is to be made for each gene, with the decision being a choice between two competing hypotheses, obtaining an estimate of some parameter of interest, or predicting the value of some function of $\mathbf{Z}_{m0}$, the observable for a new unit.

In essence, these “large M , small n ” data sets are the inputs in multiple decision problems, called in [12] as parallel inference problems, with the most common type being that of

multiple hypotheses testing. In the latter problem, for the m th gene, there is a null hypothesis H_{m0} and an alternative hypothesis H_{m1} for which a choice is to be made based on DATA. These types of problems have spurred considerable research activity among researchers, notably statisticians, since in performing multiple decision-making there is a crucial need to be cognizant and cautious of the *Hyde*-ian nature of multiplicity, though other procedures, especially those with an empirical Bayes flavor [12], have exploited the *Jekyll*-ian potentials of multiplicity [47]. In multiple hypotheses testing this entails holding a tenuous balance between two competing desires: control the rate at which correct null hypotheses are erroneously rejected, but maintain the ability to discover correct alternative hypotheses.

Similar to classical single-pair hypothesis testing, an error committed when a correct null hypothesis is rejected is referred to as of Type I, while one committed when a false null hypothesis is not rejected is of Type II. There are several types of Type I error rates in the multiple testing scenario; see, for example, [8] and [9]. In this paper we concern ourselves with the weak family wise error rate (FWER), which is the probability of rejecting at least one null hypothesis when all the nulls are correct; the strong FWER, which is the probability of rejecting at least one correct null hypothesis; and the false discovery rate (FDR), introduced by [45] and [1], which is the expected proportion of the number of false rejections of nulls relative to the number of rejections. We follow the usual convention where rejection of a null hypothesis is called a *discovery*. On the other hand, the Type II error rate of interest will be the missed discovery rate (MDR), which is the expected number of false non-rejections of nulls. There are other Type II error rates in the multiple testing setting that have been considered in the literature; see, for instance, [8], [49], [7], [11], and [9]. We justify our focus on the MDR in Section 2.

Analogous to classical hypothesis testing, in multiple hypotheses testing the commission of a Type I error is considered more serious than that of a Type II error. Therefore, one framework in the development of multiple decision functions requires that one control a chosen Type I error rate at a pre-specified level, while making the MDR, or another Type II error rate, small, possibly minimal. For example, a procedure that controls the weak FWER, under an independence assumption among the genes, is the Šidák procedure [43]; while a more conservative one, but which does not require the independence condition, is the Bonferroni procedure [4]. For control of the FDR, the procedure introduced by Benjamini and Hochberg in their seminal paper [1], hence referred to as the BH procedure, achieves the desired control. Other works have dealt with related Type I error measures to the FDR. The papers [14], [10, 11], and [53] discussed controlling the mFDR, an error rate that is asymptotically equivalent to the FDR in some settings; see [17]. On the otherhand, [48, 49] dealt with the pFDR, also similar to the FDR and related to the local FDR in [12], the latter having a Bayesian justification. Some other papers, such as [41], [27], and [24], focused on the estimation of the proportion of correct null hypotheses.

Many multiple hypotheses testing procedures, such as the Šidák, Bonferroni, and BH procedures, rely on the set of p -values of the individual tests. The validity of these p -value

based multiple testing procedures is anchored on the technical requirement that each p -value statistic is stochastically equal to or greater than a standard uniform variable under the null hypothesis. This requirement, however, is not satisfied with non-continuous variables, or when nonparametric tests, such as the Mann-Whitney-Wilcoxon two-sample test, are used. It is also not apparent whether such p -value based multiple testing procedures are utilizing, if at all, the power functions of the M tests, since many of them sets a single threshold, and genes whose p -values are smaller than this threshold are declared discovered. This approach may be acceptable in exchangeable settings, but perhaps not in situations where genes or subclasses of genes have different structures. See, for example, the papers [13] where there are subclasses of the genes possessing different structures, [15] where external covariates lead to non-exchangeability, and [34] which deals with settings with heterogeneous p -value distributions. These are situations where individual tests have different powers. Since power functions are germane for control of Type II error, would it not be the case that by imposing a common threshold on the p -values, the ability to control the Type II error rate will be compromised?

These are the motivating issues and questions for this paper. We examine these issues in a decision-theoretic framework allowing for general data types and structures enabling results to be applicable even for discrete or mixed data and with rank-based nonparametric tests. We exploit the power functions of the individual tests to develop optimal or improved procedures that control, in weak and strong senses, FWER or FDR. The procedures also possess smaller Type II error rates. We surmise that it is more the rule, rather than the exception, that in multiple hypotheses testing, the individual tests will have different power traits, owing to varied distributional characteristics among the X_m s. This could be due to differing variabilities of observed variables, differing effect sizes of interest, and possibly the use of different tests as dictated by the data type or structure, such as when some of the tests are t -tests, chi-square tests, analysis-of-variance F -tests, or nonparametric tests. It could be that the usual assumption of exchangeability of the M genes is more untenable than tenable. As a consequence, multiple testing procedures relying on the usual p -value statistics with a single-threshold rejection rule, such as the Šidák, Bonferroni, and BH procedures, cannot be expected to exploit differences in the powers of the individual tests.

There are papers dealing with multiple testing procedures which improve on single-thresholding procedures. Spjøtvoll [46], dealing with simple null and alternative hypotheses, maximized the average power among the M tests subject to a constraint on the expected number of false discoveries. Westfall et al. [58] maximized the power in replicated clinical trials involving multiple endpoints, where an adjustment is made on the significance levels of the individual tests for each of the multiple endpoints under an FWER constraint. Since the optimal solution relies on knowledge of the noncentrality parameters, a Bayesian approach was used to get a handle on these noncentrality parameters. There were also papers that approached the problem through the notion of weighted p -values such as [18], [55], [36], [25], and [34]. In some of these papers, the optimal weights of the p -values were estimated with the

aid of prior information about the distributional parameters under the alternatives. Interesting approaches toward the search for optimal multiple hypotheses testing procedures were those in Storey [50] and Storey et al. [51], where a Neyman-Pearson approach was invoked to obtain compound procedures, and Sun and Cai [53], where oracle and adaptive compound rules were developed. Compound procedures are characterized by information borrowing from each of the genes, so a decision function for a specific gene will utilize information from other genes. Decision-theoretic and Bayesian approaches were also implemented in [31], [39], [42], [12], and [19]. More recently, Efron [13] called for separate subclass analysis, while Ferkingstad, et al. [15] proposed the use of external covariates, with these papers employing Bayes and empirical Bayes approaches.

In their pioneering work, Neyman and Pearson [32] demonstrated that the most basic and fundamental type of hypotheses in the single-pair testing problem is that with simple null and alternative hypotheses. Their Fundamental Lemma, which revealed the existence and uniqueness of a most powerful (MP) test function, opened the doors to optimal classes of test functions in more complicated settings, leading to classes of test functions possessing properties such as uniformly most powerful (UMP), UMP unbiased, or UMP invariant, and led to the exploitation of the monotone likelihood ratio (MLR) property. Lehmann [28] provides a comprehensive account of the Neyman-Pearson framework of hypothesis testing, a framework which dictates that in the search for optimal test functions, the role of the power function is central and paramount. This framework also led to the divorce from the purely significance or p -value based approach to hypothesis testing which was then dominant during the first quarter of the 20th century.

It appears that, in a parallel manner, we are in the same juncture for the multiple hypotheses testing problem as almost a century ago. Many current multiple testing procedures are p -value based and do *not* exploit the power functions of the individual tests. It behooves to examine if better multiple testing procedures will arise by utilizing the individual power functions, in parallel to what Neyman and Pearson did in the single-pair hypothesis testing problem. This paper is an attempt in this direction. By considering the most basic, but we consider the most fundamental, setting in this multiple hypotheses testing situation, we will study multiple decision functions in situations where, for each gene, the null and alternative hypotheses are simple. This is also the setting considered recently in Roquian and van de Wiel [34]. In the search for multiple decision functions this will allow as starting point the most powerful test for each of the M pairs of hypotheses, with the test's existence guaranteed by the Neyman-Pearson Lemma. Each of these MP tests will have a power, but we will see that it is beneficial to look at each of these powers as functions of their MP test's size. These functions are the so-called receiver operating characteristic (ROC) functions.

We outline the contents of this paper. In Section 2 we present the decision-theoretic framework which will serve as a platform for obtaining the multiple decision functions. This entails describing the probability models, an independence condition underlying the model, relevant loss functions, multiple decision functions, and risk functions. The Type I and

Type II error rates will be informed by the choice of loss functions. We also justify the choice of the MDR as the Type II error rate of interest. Section 3 provides a review and re-examination of most powerful tests and p -values. It will also examine properties of the ROC function, which will become central in later developments. We utilize the results of Section 3 to find the optimal weak FWER-controlling procedure in Section 4. The existence will be addressed in subsection 4.2, whereas subsection 4.3 will deal with the uniqueness. Subsection 4.4 provides an explicit method for determining the optimal solution when the ROC functions are differentiable. Subsection 4.5 illustrates the theory for a specific concrete multiple testing situation involving normal distributions. In this example we show the gain in efficiency of the optimal weak FWER-controlling procedure relative to the Šidák procedure. Subsection 4.6 discusses an interesting feature of the weak FWER-controlling procedure which bears on how to optimally invest the overall FWER-size to each of the tests and point out a tangential manifestation of the strategy during the 2008 Presidential election in the United States. Section 5 discusses limitations, extensions, and connections of the problem considered. In subsection 5.1 the restriction of the optimization problem to the class of simple procedures is discussed in relation to those in [50, 46]. In subsection 5.2 extensions to situations with composite hypotheses when a monotone likelihood ratio property holds is indicated. The role of effect sizes is also discussed and some strategies are indicated when the alternative hypothesis probability measures or effect sizes are not known. Subsection 5.3 relates the weak FWER-controlling optimal procedure to p -value based procedures and distributions of p -value statistics.

Section 6 develops an improved procedure which strongly controls the FWER, whereas Section 7 develops an improved procedure which strongly controls the FDR. These new procedures possess better Type II error rate performance than the sequential Šidák and BH procedures. The development of these new procedures is anchored on the weak FWER-controlling optimal procedure in the earlier sections. It is shown that the sequential Šidák and BH procedures are special cases of these more general procedures.

In Section 8 we provide a modest simulation study demonstrating that the new FDR-controlling procedure improves on the BH procedure with respect to the MDR for the normal model considered in the simulation. Concluding remarks are provided in Section 9.

2 Mathematical Setting

Let $(\Omega, \mathcal{F}, \mathbf{P})$ be the basic probability space on which all random entities are defined and $\mathcal{M} = \{1, 2, \dots, M\}$ be an index set, with M a known positive integer. For each $m \in \mathcal{M}$, let $X_m : (\Omega, \mathcal{F}) \rightarrow (\mathcal{X}_m, \mathcal{B}_m)$, where $\mathcal{X}_m$ is some space with associated σ -field of subsets $\mathcal{B}_m$. Form the product space $(\mathcal{X}, \mathcal{B})$ with $\mathcal{X} = \times_{m \in \mathcal{M}} \mathcal{X}_m$ and $\mathcal{B} = \sigma(\times_{m \in \mathcal{M}} \mathcal{B}_m)$, so that

$$X = (X_1, X_2, \dots, X_M) : (\Omega, \mathcal{F}) \rightarrow (\mathcal{X}, \mathcal{B}).$$

The induced probability measure of X is $Q = \mathbf{P}X^{-1}$, while the (marginal) probability measure of X_m is $Q_m = PX_m^{-1}$, which is also

$$Q_m(B_m) = Q(\mathcal{X}_1 \times \dots \times \mathcal{X}_{m-1} \times B_m \times \mathcal{X}_{m+1} \times \dots \times \mathcal{X}_M), \quad \forall B_m \in \mathcal{B}_m.$$

For each $m \in \mathcal{M}$, let Q_{m0} and Q_{m1} be two known probability measures on $(\mathcal{X}_m, \mathcal{B}_m)$. In conjunction with first treating the case with simple null and alternative hypotheses for each $m \in \mathcal{M}$, we assume that Q belongs to $\mathcal{Q}$, the collection of all probability measures on $(\mathcal{X}, \mathcal{B})$ whose marginal probability measures Q_m s satisfy $Q_m \in \{Q_{m0}, Q_{m1}\}$ for each $m \in \mathcal{M}$. Let

$$\theta = (\theta_1, \dots, \theta_M) : \mathcal{Q} \rightarrow \Theta \equiv \{0, 1\}^M$$

be defined according to $\theta_m(Q) = I\{Q_m = Q_{m1}\}$, where $I\{\cdot\}$ is the indicator function. The vector $\theta(Q)$ is the state of the marginal probability measures of Q . Define, for each $Q \in \mathcal{Q}$, the subcollections

$$\mathcal{M}_0 \equiv \mathcal{M}_0(Q) = \{m \in \mathcal{M} : \theta_m(Q) = 0\}; \quad (2.1)$$

$$\mathcal{M}_1 \equiv \mathcal{M}_1(Q) = \{m \in \mathcal{M} : \theta_m(Q) = 1\}. \quad (2.2)$$

In this paper we shall impose an *independence condition* given by:

Condition (I): $(X_m, m \in \mathcal{M}_0(Q))$ is an independent collection of random entities, that is, $\forall B_m \in \mathcal{B}_m$,

$$Q(\times_{m \in \mathcal{M}_0(Q)} B_m) = \prod_{m \in \mathcal{M}_0(Q)} Q_m(B_m). \quad (2.3)$$

The collection $(X_m, m \in \mathcal{M}_1(Q))$ need not be an independent collection, but we do assume that this collection is independent of $(X_m, m \in \mathcal{M}_0(Q))$. Two extreme subcollections of $\mathcal{Q}$ are

$$\mathcal{Q}_0 = \{Q \in \mathcal{Q} : \theta_m(Q) = 0, \forall m \in \mathcal{M}\}; \quad (2.4)$$

$$\mathcal{Q}_1 = \{Q \in \mathcal{Q} : \theta_m(Q) = 1, \forall m \in \mathcal{M}\}. \quad (2.5)$$

By condition (I), $\mathcal{Q}_0$ is a singleton set. We denote by Q_0 its element. On the otherhand, $\mathcal{Q}_1$ need not be a singleton set.

Stated in its most basic form, the decision problem is to determine $\mathcal{M}_0(Q)$ and $\mathcal{M}_1(Q)$ based on X . This can be restated as a multiple hypotheses testing problem where one is interested in simultaneously testing, based on X , the M pairs of hypotheses $H_{m0} : Q_m = Q_{m0}$ versus $H_{m1} : Q_m = Q_{m1}$ for $m \in \mathcal{M}$. These pairs of hypotheses could also be stated in terms of the θ -vector via $H_{m0} : \theta_m(Q) = 0$ versus $H_{m1} : \theta_m(Q) = 1$.

We approach this problem in a decision-theoretic framework which is somewhat similar to that in Sarkar, Zhou and Ghosh [39]. The elements of this framework are as follows.

The *action space* is $\mathcal{A} = \{0, 1\}^M$ with generic element $a = (a_1, a_2, \dots, a_M)^t \in \mathcal{A}$ with the interpretation that $a_m = 0$ (1) means that H_{m0} is accepted (rejected). The *parameter space* is $\mathcal{Q}$, though the effective parameter space is $\Theta = \{0, 1\}^M$ with generic element $\theta = (\theta_1, \theta_2, \dots, \theta_M)^t$. For this decision problem, we introduce several *loss functions*, $L : \mathcal{A} \times \mathcal{Q} \rightarrow \mathbb{R}_+$, defined via

$$L_{0k}(a, Q) = I\{a^t(1 - \theta(Q)) \geq k\}, \quad k = 1, 2, \dots, M; \quad (2.6)$$

$$L_1(a, Q) = \left[\frac{a^t(1 - \theta(Q))}{a^t 1} \right] I\{a^t 1 > 0\}; \quad (2.7)$$

$$L_2(a, Q) = (1 - a)^t \theta(Q), \quad (2.8)$$

with the convention that $0/0 = 0$ and 1 is an $M \times 1$ vector of 1s. The interpretations of these loss functions are as follows. The loss function $L_{0k}(a, Q)$ equals 1 if and only if at least k false discoveries are committed, so when $k = 1$, $L_{01}(a, Q)$ becomes 1 if and only if at least one false discovery is committed. The loss $L_1(a, Q)$ could be interpreted as the *false discovery proportion*, since it is the ratio of the number of false discoveries and the number of discoveries; whereas the loss $L_2(a, Q)$ is the *number of missed discoveries* since it is the number of true alternative hypotheses that were not discovered. We focus on this missed discovery number since the relevant question is *how many correct alternatives* $(\theta(Q)^t 1)$ *were missed by using the action* a ? See also [34] which essentially uses this loss function to induce their power metric. Other types of losses, such as the false negative proportion with

$$(a, Q) \mapsto \frac{(1 - a)^t \theta(Q)}{(1 - a)^t 1} I\{(1 - a)^t 1 > 0\},$$

have also been considered in the literature such as in [17] and [39].

A *nonrandomized* multiple decision function (MDF) is a $\delta : (\mathcal{X}, \mathcal{B}) \rightarrow (\mathcal{A}, \sigma(\mathcal{A}))$, where $\sigma(\mathcal{A})$ is the power set of $\mathcal{A}$. Such an MDF may be represented by

$$\delta(x) = (\delta_1(x), \delta_2(x), \dots, \delta_M(x))^t,$$

where $\delta_m(x) \in \{0, 1\}$. In general, each δ_m could be made to depend on the full data x instead of just x_m . We denote by $\mathcal{D}$ the class of all nonrandomized MDFs. A *randomized* MDF may also be considered. Denote by $\mathcal{P}(\mathcal{A})$ the space of all probability measures over $(\mathcal{A}, \sigma(\mathcal{A}))$. A randomized MDF is a $\delta^* : (\mathcal{X}, \mathcal{B}) \rightarrow (\mathcal{P}(\mathcal{A}), \sigma(\mathcal{P}(\mathcal{A})))$. For a realization $X = x$, an action is chosen from $\mathcal{A}$ according to the probability measure $\delta^*(x)$. Denote by $\mathcal{D}^*$ the space of all possible randomized MDFs. Clearly, $\mathcal{D} \subset \mathcal{D}^*$. It is easy to see that by augmenting the data X with a randomizer $U \sim U(0, 1)$, which is independent of X , randomized MDFs could be made into nonrandomized MDFs with respect to the *augmented data* (X, U) . Henceforth, $\mathcal{D}$ will represent all *nonrandomized* MDFs $\delta(X, U)$ s based on (X, U) .

For brevity of notation, in the sequel, $\mathbf{P}_Q\{f(X, U) \in B\}$ and $E_Q\{f(X, U)\}$ will represent probability and expectation with respect to (X, U) when $X \sim Q$, $U \sim U(0, 1)$, and X and

U are independent. For each $\delta \in \mathcal{D}$ and the loss functions defined earlier, we then have the *risk functions*

$$R_{0k}(\delta, Q) = E_Q \{L_{0k}(\delta(X, U), Q)\}, k = 1, 2, \dots, M; \quad (2.9)$$

$$R_1(\delta, Q) = E_Q \{L_1(\delta(X, U), Q)\}; \quad (2.10)$$

$$R_2(\delta, Q) = E_Q \{L_2(\delta(X, U), Q)\}. \quad (2.11)$$

Given a $\delta = (\delta_1, \delta_2, \dots, \delta_M)^t$, let $\pi_\delta(Q) = (\pi_{\delta_1}(Q), \pi_{\delta_2}(Q), \dots, \pi_{\delta_M}(Q))^t$, where $\pi_{\delta_m}(Q) = E_Q\{\delta_m(X, U)\}$, be its associated vector of power functions. We may then re-express (2.11) via

$$R_2(\delta, Q) = (1 - \pi_\delta(Q))^t \theta(Q). \quad (2.12)$$

In terms of these risk functions, for an MDF $\delta \in \mathcal{D}$, its weak FWER is $\text{FWER}(\delta) = R_{01}(\delta, Q_0)$, and its weak k -FWER is $k\text{-FWER}(\delta) = R_{0k}(\delta, Q_0)$ for $k \geq 1$. If each δ_m depends only on X_m and U , by condition (I),

$$\text{FWER}(\delta) = 1 - E \left\{ \prod_{m \in \mathcal{M}} [1 - \mathbf{P}_{Q_{m0}}\{\delta_m(X_m, U) = 1 | U\}] \right\} \quad (2.13)$$

where expectation is over the randomizer U . An alternative formulation when $Q = Q_0$ and with the m th component δ_m^* of the randomized MDF δ^* depending only on X_m is to have $U = (U_1, U_2, \dots, U_M)$ consisting of independent and identically distributed $U(0, 1)$ variables and independent of the X_m s. The m th component $\delta_m^*(X_m) (\in [0, 1])$ may be redefined via $\delta_m(X_m, U_m) = I\{U_m \leq \delta_m^*(X_m)\}$, making $\delta(X, U) = (\delta_m(X_m, U_m), m \in \mathcal{M})^t$ a nonrandomized MDF depending on (X, U) . In this case, (2.13) becomes

$$\text{FWER}(\delta) = 1 - \prod_{m \in \mathcal{M}} [1 - \mathbf{P}_{Q_{m0}}\{\delta_m(X_m, U_m) = 1\}]. \quad (2.14)$$

The risk function $R_1(\delta, Q)$ is the false discovery *rate* (FDR) of δ at Q , the error rate introduced in [1]; while the risk function $R_2(\delta, Q)$ will be called the missed discovery *rate* (MDR) of δ at Q . The adjective ‘rate’ may be somewhat misleading since $R_2(\delta, Q)$ takes values in $[0, |\mathcal{M}_1(Q)|]$ instead of $[0, 1]$; however, this does not cause difficulty since given the true underlying probability measure Q of X , $|\mathcal{M}_1(Q)|$ is a constant. This risk is related to the expected number of true positives (ETP), an error measure used in [46] and [50], via $\text{ETP}(\delta, Q) = |\mathcal{M}_1(Q)| - R_2(\delta, Q)$; see subsection 5.1 for more discussions of these error measures.

In analogy with the Neyman-Pearson framework for testing a null hypothesis versus an alternative hypothesis, the risks $R_{0k}(\delta, Q)$ and $R_1(\delta, Q)$ will be viewed as Type I error rates, whereas the risk $R_2(\delta, Q)$ will be a Type II error rate. As Type I errors are considered more serious than Type II errors, in the search for good MDFs, Type I error rates are required

not to exceed a pre-specified threshold. Subject to this constraint, an MDF is to be chosen whose Type II error rate is small, if not minimal.

Thus, to find an optimal MDF that provides *weak* FWER control in a subclass $\mathcal{D}_0 \subseteq \mathcal{D}$, a threshold $\alpha \in (0, 1)$ is specified, and it is desired to find an MDF $\delta^* \in \mathcal{D}_0$ with $R_{01}(\delta^*, Q_0) = \text{FWER}(\delta^*) \leq \alpha$, and such that for any other $\delta \in \mathcal{D}_0$ satisfying $R_{01}(\delta, Q_0) = \text{FWER}(\delta) \leq \alpha$, we have

$$\sup_{Q \in \mathcal{Q}} R_2(\delta^*, Q) \leq \sup_{Q \in \mathcal{Q}} R_2(\delta, Q). \quad (2.15)$$

The criterion in (2.15) is of a minimax flavor. One may require only that $R_2(\delta^*, Q^*) \leq R_2(\delta, Q^*)$ where Q^* is the true, but unknown, probability law of X ; however, this condition may be too strong so as to preclude a solution to the optimization problem, though see Storey [50] (also Section 5) for a situation which uses a different Type I error and where an optimal, albeit an oracle, solution with respect to minimizing $R_2(\delta, Q^*)$ is possible. Observe that for any $\delta \in \mathcal{D}$, using the representation of $R_2(\delta, Q)$ in (2.12),

$$\sup_{Q \in \mathcal{Q}} R_2(\delta, Q) = \sup_{Q \in \mathcal{Q}_1} R_2(\delta, Q) = M - \inf_{Q \in \mathcal{Q}_1} \sum_{m \in \mathcal{M}} \pi_{\delta_m}(Q). \quad (2.16)$$

Thus, the optimality condition on the MDR in (2.15) is equivalent to maximizing the sum of the powers of the individual components of the MDF, that is, maximizing $\sum_{m \in \mathcal{M}} \pi_{\delta_m}(Q_{m1})$. Interestingly, if we had innocently standardized the loss function $L_2(a, Q)$ in order to take values in $[0, 1]$ via division by $|\mathcal{M}_1(Q)| = \theta(Q)^t 1$, the number of correct alternatives, then the minimax justification above does not carry through!

In contrast, for *strong* FWER control, given the threshold of α , it is desired to find a compound MDF, $\delta^* \in \mathcal{D}$, such that

$$R_{01}(\delta^*, Q^*) \leq \alpha \quad (2.17)$$

whatever the true, but unknown, probability law Q^* of X is, and such that $\sum_{m \in \mathcal{M}} \pi_{\delta_m^*}(Q_{m1})$ is large, if not maximal, among all $\delta \in \mathcal{D}$ satisfying $R_{01}(\delta, Q^*) \leq \alpha$. Clearly, analogous requirements exist for weak and strong k -FWER control. However, in this paper, due to space limitations, we will not focus on this k -FWER error rate for $k > 1$.

For (strong) FDR-control, a threshold $q^* \in (0, 1)$ is specified and we seek a compound MDF, $\delta^* \in \mathcal{D}$, such that, *whatever* Q^* is,

$$R_1(\delta^*, Q^*) \leq q^*, \quad (2.18)$$

and for any other $\delta \in \mathcal{D}$ satisfying $R_1(\delta, Q^*) \leq q^*$, we have

$$\inf_{Q \in \mathcal{Q}_1} \sum_{m \in \mathcal{M}} \pi_{\delta_m^*}(Q) \geq \inf_{Q \in \mathcal{Q}_1} \sum_{m \in \mathcal{M}} \pi_{\delta_m}(Q).$$

The crucial aspect in strong control, both for FWER and FDR, is that the constraints in (2.17) and (2.18) need to hold *whatever* the unknown Q^* is, in contrast to the weak FWER constraint where it is only required at $Q_0 \in \mathcal{Q}_0$. For more discussion of weak and strong control, see [8] and [9], and for discussion of optimality conditions in multiple testing, refer to [29], wherein maximin optimality results were established for some step-down and step-up MTPs.

3 Revisiting MP Tests and p -Value Statistics

The initial subclass of $\mathcal{D}$ which is of interest is $\mathcal{D}_0$, the subclass of *simple* MDFs. A simple MDF $\delta(X, U)$ is one whose m th component δ_m depends only on (X_m, U_m) for every $m \in \mathcal{M}$. Within $\mathcal{D}_0$ we examine the existence, uniqueness, and structure of an optimal MDF that controls FWER in the weak sense. This optimal MDF will serve later as anchor to developing improved MDFs that strongly control FWER and FDR. These new MDFs will be outside the subclass $\mathcal{D}_0$, hence are called *compound* MDFs.

But, first, for the single-pair hypotheses testing problem, we introduce the notion of a test or decision process together with its ROC function and p -value statistic. We revisit most powerful (MP) tests and obtain properties of the ROC function and the p -value statistic associated with the MP test process. These properties will be needed in later developments.

3.1 Decision Processes and ROC Functions

Let $X : (\Omega, \mathcal{A}) \rightarrow (\mathcal{X}, \mathcal{B})$ be an observable random entity and $Q = \mathbf{P}X^{-1}$ its probability measure. Consider testing, based on X , the pair of hypotheses $H_0 : Q = Q_0$ versus $H_1 : Q = Q_1$, where Q_0 and Q_1 are two probability measures on $(\mathcal{X}, \mathcal{B})$. Denote by q_0 and q_1 versions of the density functions of Q_0 and Q_1 , respectively, with respect to some fixed dominating measure ν , e.g., $\nu = Q_0 + Q_1$, or it could be counting or Lebesgue measure. Recall that a test function is a measurable function $\delta : (\mathcal{X}, \mathcal{B}) \rightarrow ([0, 1], \sigma[0, 1])$, where $\sigma[0, 1]$ is the Borel sigma-field on $[0, 1]$. Given $X = x$, $\delta(x)$ is the probability of deciding in favor of H_1 . Its size is $\alpha_\delta = E_{Q_0}\delta(X)$. It is of level $\alpha \in [0, 1]$, if $\alpha_\delta \leq \alpha$. Its power is $\pi_\delta = E_{Q_1}\delta(X)$. Recall that a test function δ^* is most powerful (MP) of level α if $\alpha_{\delta^*} \leq \alpha$ and for any other test function δ with $\alpha_\delta \leq \alpha$, we have $\pi_{\delta^*} \geq \pi_\delta$. We now introduce the notion of a decision process and its associated ROC curve.

Definition 3.1 A collection $\Delta = \{\delta_\eta : \eta \in [0, 1]\}$ of test functions satisfying the conditions that, a.e. $[Q]$, $\delta_0(x) = 0$, $\delta_1(x) = 1$, and $\eta \mapsto \delta_\eta(x)$ is nondecreasing and right-continuous, is a decision process.

Definition 3.2 For a decision process $\Delta = \{\delta_\eta : \eta \in [0, 1]\}$, its size function is $A_\Delta : [0, 1] \rightarrow [0, 1]$ and its power function is $\rho_\Delta : [0, 1] \rightarrow [0, 1]$, where $A_\Delta(\eta) = \alpha_{\delta_\eta} = E_{Q_0}\delta_\eta(X)$ and

$\rho_\Delta(\eta) = \pi_{\delta_\eta} = E_{Q_1}\delta_\eta(X)$. Its receiver operating characteristic (ROC) curve is $ROC(\Delta) \equiv \text{Graph}\{(A_\Delta(\eta), \rho_\Delta(\eta)) : \eta \in [0, 1]\}$. When $A_\Delta(\eta) = \eta$ for all $\eta \in [0, 1]$, the mapping $\eta \mapsto \rho_\Delta(\eta)$ is called the ROC function of Δ .

The use of the phrase *power function* in Definition 3.2 may be construed a misnomer since we are not viewing this as a function of a parameter as is the usual meaning of the phrase. However, for lack of a better name, we adopt this terminology. Also, when the decision process Δ satisfies the condition that $\forall \eta \in [0, 1] : A_\Delta(\eta) = \eta$, it is said to be *size-valid*. In the sequel, when convenient notationally, δ_η and $\delta(\eta)$ will be used interchangeably to represent $\delta(\cdot; \eta)$.

Let $L : (\mathcal{X}, \mathcal{B}) \rightarrow (\mathfrak{R}_+, \sigma(\mathfrak{R}_+))$ be a version of the likelihood ratio function, so $L(x) = q_1(x)/q_0(x)$ a.e. $[\nu]$. Let $G_0(\cdot)$ and $G_1(\cdot)$ be the distribution functions of $L(X)$ when $\mathcal{L}(X) = Q_0$ and $\mathcal{L}(X) = Q_1$, respectively, where $\mathcal{L}(X)$ is probability measure of X . For a monotone nondecreasing right-continuous function $M(\cdot)$ from $\mathfrak{R}$ into $\mathfrak{R}$, let

$$M^{-1}(r) = \inf\{x \in \mathfrak{R} : M(x) \geq r\} \quad \text{and} \quad \Delta M(r) = M(r) - M(r-).$$

The Neyman-Pearson Fundamental Lemma [32] states that the MP test function of level η for testing H_0 versus H_1 is

$$\delta^*(X; \eta) \equiv \delta_\eta^* = I\{L(X) > c(\eta)\} + \gamma(\eta)I\{L(X) = c(\eta)\}, \quad (3.1)$$

where $c(\eta) = G_0^{-1}(1 - \eta)$ and $\gamma(\eta) = (G_0(c(\eta)) - (1 - \eta))/\Delta G_0(c(\eta))$. Let $U \sim U(0, 1)$ be independent of X . Redefine δ^* via

$$\delta^{**}(X, U; \eta) = \delta_\eta^{**} = I\{\delta^*(X; \eta) = 1\} + I\{\delta^*(X; \eta) = \gamma(\eta); U \leq \gamma(\eta)\}.$$

This is nonrandomized with respect to (X, U) , so with the aid of the auxiliary randomizer U , the MP test could always be made nonrandomized. The decision process formed from these MP tests, given by

$$\Delta^* = \{\delta_\eta^* : \eta \in [0, 1]\} = \{\delta_\eta^{**} : \eta \in [0, 1]\}, \quad (3.2)$$

is referred to as the most powerful (MP) decision process. The power (at $Q = Q_1$) of the MP test δ_η^* or δ_η^{**} is

$$\rho_{\Delta^*}(\eta) \equiv \pi_{\delta_\eta^*} = \pi_{\delta_\eta^{**}} = 1 - G_1(c(\eta)) + \gamma(\eta)\Delta G_1(c(\eta)). \quad (3.3)$$

It is well-known (see [28]) that if $\pi_{\delta_\eta^*} < 1$, then $\alpha_{\delta_\eta^*} = \eta$. We denote by A_{Δ^*} and ρ_{Δ^*} the size and power functions, respectively, of Δ^* . Note that if $\pi_{\delta_\eta^*} < 1$ for all $\eta < 1$, then the mapping $\eta \mapsto \rho_{\Delta^*}(\eta)$ is the ROC function of the MP decision process.

Proposition 3.1 *The function $\rho_{\Delta^*} : [0, 1] \rightarrow [0, 1]$ in (3.3) satisfies $\rho_{\Delta^*}(\eta) \geq \eta$, concave, continuous, and nondecreasing. Furthermore, it is strictly increasing on the set $\mathcal{N}_{<} \equiv \{\eta \in [0, 1] : \rho_{\Delta^*}(\eta) < 1\}$.*

Proof: Recall that $\rho_{\Delta^*}(\eta) \equiv \pi_{\delta^*(\eta)}$. The first result follows from the unbiasedness property of MP tests. Suppose $\rho_{\Delta^*}(\cdot)$ is not concave. Then there must exist $\eta_1 \in [0, 1]$, $\eta_2 \in [0, 1]$, and $\xi \in (0, 1)$ such that

$$\xi \pi_{\delta^*(\eta_1)} + (1 - \xi) \pi_{\delta^*(\eta_2)} > \pi_{\delta^*(\xi \eta_1 + (1 - \xi) \eta_2)}. \quad (3.4)$$

Consider the test function $\delta^{**} = \xi \delta^*(\eta_1) + (1 - \xi) \delta^*(\eta_2)$. The size of this test is $\xi \eta_1 + (1 - \xi) \eta_2$, while its power is $\xi \pi_{\delta^*(\eta_1)} + (1 - \xi) \pi_{\delta^*(\eta_2)}$. From (3.4) the power of δ^{**} exceeds that of the MP test whose size is $\xi \eta_1 + (1 - \xi) \eta_2$. Since δ^{**} has the same size as this MP test, this is a contradiction. Thus, $\eta \mapsto \pi_{\delta^*(\eta)}$ must be concave, and hence continuous. Furthermore, since $\eta \leq \pi_{\delta^*(\eta)}$ with $\pi_{\delta^*(1)} = 1$, it follows by concavity that it is nondecreasing.

Suppose that on $\mathcal{N}_{<}$, $\rho_{\Delta^*}(\cdot)$ is *not* strictly increasing. Then there exists an $\eta_0 \in \mathcal{N}_{<}$ such that for some $\epsilon > 0$ and $r < 1$, $\rho_{\Delta^*}(\eta) = r$ whenever $\eta \in [\eta_0 - \epsilon, \eta_0] \subset [0, 1]$. Let $\eta_1 = \sup\{\eta \in [0, 1] : \rho_{\Delta^*}(\eta) = r\}$. Continuity and nondecreasingness of $\rho_{\Delta^*}(\cdot)$ imply that $\eta_1 < 1$. Furthermore, since $\rho_{\Delta^*}(1) = 1$, then there exists an $\eta_2 \in (\eta_1, 1]$ with $1 \geq \rho_{\Delta^*}(\eta_2) > r$. Since $r = \rho_{\Delta^*}(\eta_1) < \frac{1}{2}[\rho_{\Delta^*}(\eta_0 - \epsilon) + \rho_{\Delta^*}(\eta_2)]$, then on $[\eta_0 - \epsilon, \eta_2]$, $\rho_{\Delta^*}(\cdot)$ is not concave. This contradicts concavity of $\rho_{\Delta^*}(\cdot)$, so the supposition could not hold. Therefore, $\rho_{\Delta^*}(\cdot)$ is strictly increasing on $\mathcal{N}_{<}$. $\parallel$

3.2 p -Value Statistics

We introduce randomized p -value statistics through the framework of decision processes.

Definition 3.3 *Let $\Delta = \{\delta_\eta : \eta \in [0, 1]\}$ be a decision process, where $\delta_\eta : (\mathcal{X}, \mathcal{B}) \rightarrow ([0, 1], \sigma[0, 1])$. Let $U \sim U[0, 1]$ and independent of X . The (randomized) p -value statistic associated with Δ is $S_\Delta : (\mathcal{X} \times [0, 1], \mathcal{B} \otimes \sigma[0, 1]) \rightarrow ([0, 1], \sigma[0, 1])$ with $S_\Delta(x, u) = \inf\{\eta \in [0, 1] : u \leq \delta_\eta(x)\}$.*

When the δ_η s in Definition 3.3 are nonrandomized, $S_\Delta(X, U)$ coincides with the usual p -value statistic. However, when the δ_η s are randomized, $S_\Delta(X, U)$ is a *randomized* p -value statistic. See also [5] for a more specialized definition of a randomized p -value statistic. Let us denote by $H_0(\cdot)$ and $H_1(\cdot)$ the distribution functions of $S_\Delta(X, U)$ when $\mathcal{L}(X) = Q_0$ and $\mathcal{L}(X) = Q_1$, respectively.

Proposition 3.2 *Let $\Delta = \{\delta_\eta : \eta \in [0, 1]\}$ be a decision process. Then, for all $s \in [0, 1]$, $H_0(s) = A_\Delta(s)$ and $H_1(s) = \pi_{\delta(s)} = \rho_\Delta(s)$.*

Proof: Using properties of a decision process, specifically the a.e.-[Q] right-continuity of $\eta \mapsto \delta_\eta(x)$, then a.e.-[Q], for each $s \in [0, 1]$, $\{(x, u) : S_\Delta(x, u) \leq s\} = \{(x, u) : u \leq \delta_s(x)\}$. Consequently, for $j = 0, 1$, and since $U \sim U[0, 1]$ and $X \perp U$, we have $H_j(s) = \mathbf{P}_{Q_j}\{S_\Delta(X, U) \leq s\} = \mathbf{P}_{Q_j}\{U \leq \delta_s(X)\} = E_{Q_j}\mathbf{P}_{Q_j}\{U \leq \delta_s(X)|U\} = E_{Q_j}\delta_s(X)$, which equals $A_\Delta(s)$ when $j = 0$ and $\rho_\Delta(s)$ when $j = 1$. $\parallel$

Corollary 3.1 *For a decision process Δ , S_Δ has a standard uniform distribution under $\mathcal{L}(X) = Q_0$ if and only if $A_\Delta(\eta) = \eta$ for all $\eta \in [0, 1]$.*

This result is immediate from Proposition 3.2. Note that the conclusion of Corollary 3.1 holds for the MP decision process Δ^* in (3.2) provided $\rho_{\Delta^*}(\eta) < 1$ for $\eta < 1$. We also highlight the last result in Proposition 3.2 which states that the ROC function $\rho_\Delta(\cdot)$ equals $H_1(\cdot)$, the distribution of the p -value statistic under the alternative hypothesis. The p -value statistic as defined in Definition 3.3 is general and applicable even with discrete or mixed data and with randomized test functions. We refer the reader to [20] for properties of this randomized p -value statistic and its use in existing FDR-controlling procedures.

4 Optimal *Weak* FWER Control

Let us now return to the multiple decision problem formulated in Section 2. We first focus our attention on the subclass $\mathcal{D}_0$ consisting of simple decision functions.

Definition 4.1 *A collection $\Delta = (\Delta_m : m \in \mathcal{M})$, where $\Delta_m = (\delta_m(\eta) : \eta \in [0, 1])$ is a decision process on $(\mathcal{X} \times [0, 1]^M, \mathcal{B} \otimes \sigma[0, 1]^M)$, is called a multiple decision process (MDP). It is simple if each Δ_m is simple; otherwise, it is compound.*

Definition 4.2 *For a simple MDP $\Delta = (\Delta_m : m \in \mathcal{M})$, its associated multiple decision size function is $\mathbf{A}_\Delta = (A_{\Delta_m} : m \in \mathcal{M})$ and its multiple decision ROC function is $\rho_\Delta = (\rho_{\Delta_m} : m \in \mathcal{M})$, where A_{Δ_m} and ρ_{Δ_m} are the size and ROC functions, respectively, of Δ_m .*

4.1 Optimization Problem

Let us suppose that we are given a simple MDP Δ . Then a multiple decision size vector $\eta = (\eta_m : m \in \mathcal{M}) \in \mathcal{N} \equiv [0, 1]^M$ will determine from Δ an MDF $\delta_\Delta(\eta) = (\delta_m(\eta_m) : m \in \mathcal{M})^t$, belonging to $\mathcal{D}_0$. For this MDF,

$$R_{01}(\delta_\Delta(\eta), Q_0) = \text{FWER}(\delta_\Delta(\eta)) = 1 - \prod_{m \in \mathcal{M}} [1 - A_{\Delta_m}(\eta_m)]$$

and, for $Q_1 \in \mathcal{Q}_1$, $R_2(\delta_{\Delta}(\eta), Q_1) = M - \sum_{m \in \mathcal{M}} \rho_{\Delta_m}(\eta_m)$. Let us fix an FWER-threshold of $\alpha \in (0, 1)$. Suppose there exists a multiple decision size vector $\eta_{\Delta}^*(\alpha) \in \mathcal{N}$ such that

$$\eta_{\Delta}^*(\alpha) = \arg \max_{\eta \in \mathcal{N}} \left\{ \sum_{m \in \mathcal{M}} \rho_{\Delta_m}(\eta_m) : \prod_{m \in \mathcal{M}} [1 - A_{\Delta_m}(\eta_m)] \geq 1 - \alpha \right\}.$$

Then, $\mathbf{A}_{\Delta}(\eta_{\Delta}^*(\alpha)) = (A_{\Delta_m}(\eta_{\Delta,m}^*(\alpha)) : m \in \mathcal{M})$ is the optimal multiple decision size vector for weak FWER control at α associated with the simple MDP Δ . The associated optimal simple MDF is $\delta_{\Delta}(\mathbf{A}_{\Delta}(\eta_{\Delta}^*(\alpha)))$.

However, since H_{m0} and H_{m1} are both simple, from Section 3.1, there exists a simple most powerful MDP, $\Delta^* = (\Delta_m^* : m \in \mathcal{M})$, where $\Delta_m^* = (\delta_m^*(\eta) : \eta \in [0, 1])$ with $\delta_m^*(\eta)$ the simple Neyman-Pearson MP test function of size η for H_{m0} versus H_{m1} . Consider the simple MDF obtained from Δ^* given by $(\delta_m^*(A_{\Delta_m}(\eta_{\Delta,m}^*(\alpha))) : m \in \mathcal{M})$. Then, this simple MDF will satisfy the FWER constraint, and by virtue of the MP property of each $\delta_m^*(A_{\Delta_m}(\eta_{\Delta,m}^*(\alpha)))$ for each $m \in \mathcal{M}$, it follows that

$$\sum_{m \in \mathcal{M}} \rho_{\Delta_m^*}(A_{\Delta_m}(\eta_{\Delta,m}^*(\alpha))) \geq \sum_{m \in \mathcal{M}} \rho_{\Delta_m}(A_{\Delta_m}(\eta_{\Delta,m}^*(\alpha))).$$

This implies that in searching for the optimal weak FWER-controlling simple MDF it suffices to restrict to the simple most powerful MDP Δ^* . Without loss of generality, we may assume $A_{\Delta_m^*}(\eta) = \eta$ for $m \in \mathcal{M}$ and $\eta \in [0, 1]$. The optimization problem amounts to obtaining a multiple decision size vector $\eta_{\Delta^*}^*(\alpha) \in \mathcal{N}$ satisfying

$$\eta_{\Delta^*}^*(\alpha) = \arg \max_{\eta \in \mathcal{N}} \left\{ \sum_{m \in \mathcal{M}} \rho_{\Delta_m^*}(\eta_m) : \prod_{m \in \mathcal{M}} (1 - \eta_m) \geq 1 - \alpha \right\}, \quad (4.1)$$

provided that such a multiple decision size vector exists. The optimal *weak* FWER-controlling simple MDF is then

$$\delta_W^*(\alpha) \equiv (\delta_m^*(\eta_{\Delta^*,m}^*(\alpha)) : m \in \mathcal{M}). \quad (4.2)$$

We mention two choices for the size vector $\eta = (\eta_m : m \in \mathcal{M})$ satisfying the FWER constraint in (4.1). The Šidák procedure [43] has

$$\eta_m = \eta_m(\alpha) = 1 - (1 - \alpha)^{1/M}, \quad m \in \mathcal{M}. \quad (4.3)$$

This guarantees that the FWER is exactly equal to α , though this requires for its validity the independence condition (I); see (2.3). A conservative choice of η is the Bonferroni inequality-derived choice with

$$\eta_m = \eta_m(\alpha) = \alpha/M, \quad m \in \mathcal{M}. \quad (4.4)$$

This choice also satisfies the FWER constraint, though equality is not achieved. However, it does not require the independence condition (I).

4.2 Existence of Optimal Size Vector

This subsection establishes the existence of an optimal multiple decision size vector for weak FWER control when dealing with $\mathcal{D}_0$. As pointed out in the preceding subsection, it suffices to look for the optimal weak FWER-controlling simple MDF by starting with the most powerful simple MDP $\Delta^* = (\Delta_m^* : m \in \mathcal{M})$. For brevity, we write $\rho_m \equiv \rho_{\Delta_m^*}$ and $A_m(\eta) \equiv A_{\Delta_m^*}(\eta) = \eta$. Recall that $\mathcal{N} = [0, 1]^M$, the multiple decision size space. For $\alpha \in [0, 1]$, define

$$C_\alpha = \begin{cases} \{\eta \in \mathcal{N} : \sum_{m \in \mathcal{M}} \log(1 - \eta_m) \geq \log(1 - \alpha)\} & \text{if } \alpha < 1 \\ \mathcal{N} & \text{if } \alpha = 1 \end{cases}, \quad (4.5)$$

the weak FWER constraint set. The following provides properties of C_α .

Proposition 4.1 *C_α satisfies (i) $\eta = \mathbf{0} \in C_\alpha$; (ii) $(\mathbf{0}, \alpha_m) \in C_\alpha$ for all $m \in \mathcal{M}$, where $(\mathbf{0}, \alpha_m)$ is the zero-vector with the m th element replaced by α ; and (iii) it is convex and closed.*

Proof: The results clearly hold when $\alpha = 1$ since $C_\alpha = \mathcal{N}$. For $\alpha \in [0, 1)$, results (i) and (ii) are immediate, while the closedness of C_α follows from the continuity of the logarithm function. Let $\eta_1, \eta_2 \in C_\alpha$ with $\eta_1 \neq \eta_2$, and let $\xi \in (0, 1)$. Since $\sum_{m \in \mathcal{M}} \log(1 - \eta_{jm}) \geq \log(1 - \alpha)$, $j = 1, 2$, and the mapping $\eta \mapsto \log(1 - \eta)$ is strictly concave, then $\sum_{m \in \mathcal{M}} \log[1 - (\xi\eta_{1m} + (1 - \xi)\eta_{2m})] > \xi \sum_{m \in \mathcal{M}} \log(1 - \eta_{1m}) + (1 - \xi) \sum_{m \in \mathcal{M}} \log(1 - \eta_{2m}) \geq \xi \log(1 - \alpha) + (1 - \xi) \log(1 - \alpha) = \log(1 - \alpha)$. This establishes the convexity of C_α . $\parallel$

Definition 4.3 *The upper set of $\eta_0 \in \mathcal{N}$ is $U(\eta_0) = \{\eta \in \mathcal{N} : \eta_m \geq \eta_{0m}, \forall m \in \mathcal{M}\}$. The upper boundary set of the constraint set C_α is $UB(C_\alpha) = \{\eta \in \mathcal{N} : C_\alpha \cap U(\eta) = \{\eta\}\}$.*

Proposition 4.2 *For all $\alpha \in [0, 1)$, $UB(C_\alpha) = \{\eta \in \mathcal{N} : \sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha)\}$.*

Proof: Let $\eta \in UB(C_\alpha)$ so $\{\eta\} = C_\alpha \cap U(\eta)$. Suppose $\sum_{m \in \mathcal{M}} \log(1 - \eta_m) > \log(1 - \alpha)$. Then by continuity of the logarithm function, there exists an $\epsilon > 0$ such that $\eta + \mathbf{1}\epsilon \in \mathcal{N}$ and $\sum_{m \in \mathcal{M}} \log(1 - \eta_m) > \sum_{m \in \mathcal{M}} \log[1 - (\eta_m + \epsilon)] \geq \log(1 - \alpha)$. Thus, $\eta + \mathbf{1}\epsilon \in C_\alpha$ and clearly $\eta + \mathbf{1}\epsilon \in U(\eta)$. Consequently, $\eta + \mathbf{1}\epsilon \in C_\alpha \cap U(\eta)$ which contradicts the fact that $\{\eta\} = C_\alpha \cap U(\eta)$. Therefore, we must have $\sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha)$.

On the other hand, let $\eta \in \mathcal{N}$ such that $\sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha)$. Then $\eta \in C_\alpha$, and since $\eta \in U(\eta)$, it follows that $\eta \in C_\alpha \cap U(\eta)$. Suppose there exists an $\eta_1 \in \mathcal{N}$ with $\eta_1 \neq \eta$ and $\eta_1 \in U(\eta)$. Then, $\eta_1 = \eta + \gamma$ with $\gamma_m \geq 0$ for all $m \in \mathcal{M}$ with strict inequality for some $m \in \mathcal{M}$. Therefore, $\sum_{m \in \mathcal{M}} \log(1 - \eta_{1m}) = \sum_{m \in \mathcal{M}} \log(1 - \eta_m - \gamma_m) < \sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha)$. This implies that $\eta_1 \notin C_\alpha$. Therefore we must have $\{\eta\} = C_\alpha \cap U(\eta)$, hence $\eta \in UB(C_\alpha)$. $\parallel$

Proposition 4.3 $\mathcal{N}_b \equiv \{\eta \in \mathcal{N} : \sum_{m \in \mathcal{M}} \rho_m(\eta_m) \geq Mb\}$ for $b \in [0, 1]$ satisfies (i) $\eta = \mathbf{1} \in \mathcal{N}_b$, (ii) it is closed and convex, and (iii) $\mathcal{N} = \mathcal{N}_0 \supseteq \mathcal{N}_{b_1} \supseteq \mathcal{N}_{b_2}$ for $0 \leq b_1 \leq b_2 \leq 1$.

Proof: From Proposition 3.1, $\forall m \in \mathcal{M}, \rho_m(1) = 1$, hence $\mathbf{1} \in \mathcal{N}_b$. It was also established in the same proposition that $\eta \mapsto \rho_m(\eta)$ is continuous, nondecreasing, and concave. That $\mathcal{N}_b$ is closed follows from the continuity of each $\rho_m(\cdot)$. The convexity of $\mathcal{N}_b$ follows from the concavity of each of the $\rho_m(\cdot)$ analogously in the proof of Proposition 4.1. The last result is immediate from the definition of $\mathcal{N}_b$. $\parallel$

Proposition 4.4 Let $B_\alpha = \{b \in [0, 1] : \mathcal{N}_b \cap C_\alpha \neq \emptyset\}$ for $\alpha \in [0, 1)$ and let $b_\alpha^* = \sup B_\alpha$. Then $B_\alpha = [0, b_\alpha^*]$.

Proof: Obviously $0 \in B_\alpha$, so B_α is nonempty, and hence b_α^* is well-defined. Let $b > 0$ with $b \in B_\alpha$. Let $b_1 \in [0, b)$. From (iii) of Proposition 4.3, $\mathcal{N}_{b_1} \supseteq \mathcal{N}_b$, hence, since $\mathcal{N}_b \cap C_\alpha \neq \emptyset$, then $\mathcal{N}_{b_1} \cap C_\alpha \neq \emptyset$. Therefore, $b_1 \in B_\alpha$. Let $\{b_n : n = 1, 2, \dots\}$ be a sequence in B_α such that $b_n \uparrow b_\alpha^*$. For each $n = 1, 2, \dots$ there exists an $\eta_n \in \mathcal{N}$ such that $\eta_n \in C_\alpha$ and $\sum_{m \in \mathcal{M}} \rho_m(\eta_{nm}) \geq Mb_n$. Consider the sequence $\{\eta_n\}$ in $\mathcal{N}$. This is a sequence belonging to the closed and bounded set C_α . By the Bolzano-Weierstrass Theorem [35], there exists a subsequence $\{\eta_{n'}\}$ of $\{\eta_n\}$ such that for some $\eta_0 \in C_\alpha$, $\eta_{n'} \rightarrow \eta_0$. Furthermore, since the $\rho_m(\cdot)$ s are continuous, then $\sum_{m \in \mathcal{M}} \rho_m(\eta_{0m}) = \lim_{n' \rightarrow \infty} \sum_{m \in \mathcal{M}} \rho_m(\eta_{n'm}) \geq M \lim_{n' \rightarrow \infty} b_{n'} = Mb_\alpha^*$. Therefore, $\eta_0 \in \mathcal{N}_{b_\alpha^*}$, hence $\eta_0 \in C_\alpha \cap \mathcal{N}_{b_\alpha^*}$. Thus, $b_\alpha^* \in B_\alpha$. $\parallel$

Theorem 4.1 (Existence) Let $\alpha \in [0, 1)$. Then $C_\alpha \cap \mathcal{N}_{b_\alpha^*} \neq \emptyset$. Furthermore, $\eta \in \mathcal{N}$ is a weak FWER- α optimal multiple decision size vector if and only if $\eta \in C_\alpha \cap \mathcal{N}_{b_\alpha^*}$.

Proof: First, observe from Proposition 4.4 that $C_\alpha \cap \mathcal{N}_{b_\alpha^*} \neq \emptyset$. Each element $\eta_0 \in C_\alpha \cap \mathcal{N}_{b_\alpha^*}$ satisfies the FWER- α constraint and also achieves the optimal (largest) value of $\sum_{m \in \mathcal{M}} \rho_m(\eta_m)$ among all $\eta \in C_\alpha$. Therefore, η_0 is an optimal size vector for FWER control at α .

Suppose that η_0 is an FWER- α optimal solution but $\eta_0 \notin C_\alpha \cap \mathcal{N}_{b_\alpha^*}$. Then there must exist a $b > b_\alpha^*$ such that $\eta_0 \in \mathcal{N}_b$. Since $\eta_0 \in C_\alpha$, then we have $C_\alpha \cap \mathcal{N}_b \neq \emptyset$. But this contradicts the maximality of b_α^* . Hence, the supposition could not be true. $\parallel$

4.3 Uniqueness of Optimal Size Vector

Theorem 4.1 guarantees the existence of an optimal weak FWER multiple decision size vector, but does not address whether the solution is unique. We consider this uniqueness issue in this subsection. For this purpose, we first define sections of C_α .

Definition 4.4 Let $\alpha \in (0, 1)$ and C_α be the constraint set. The m th section of C_α is the subset of $[0, 1]$ given by $C_\alpha(m) = \{\eta_m \in [0, 1] : \eta \in C_\alpha\}$.

Theorem 4.2 (Uniqueness) *Let $\alpha \in [0, 1)$. If, $\forall m \in \mathcal{M}, \eta_m \mapsto \rho_m(\eta_m)$ is strictly increasing on $C_\alpha(m)$, then the optimal weak FWER- α multiple decision size vector is unique and is the η^* with $C_\alpha \cap \mathcal{N}_{b_\alpha^*} = \{\eta^*\}$.*

Proof: It suffices to show from Theorem 4.1 that $C_\alpha \cap \mathcal{N}_{b_\alpha^*}$ is a singleton set. Suppose it is not a singleton set. Let $\eta_1, \eta_2 \in \mathcal{N}$ with $\eta_1 \neq \eta_2$ such that for $j = 1, 2$, $\sum_{m \in \mathcal{M}} \log(1 - \eta_{jm}) \geq \log(1 - \alpha)$ and $\sum_{m \in \mathcal{M}} \rho_m(\eta_{jm}) \geq Mb_\alpha^*$. Let $\xi \in (0, 1)$ and define $\eta^* = \xi\eta_1 + (1 - \xi)\eta_2$. By convexity of both C_α and $\mathcal{N}_{b_\alpha^*}$, we have $\eta^* \in C_\alpha \cap \mathcal{N}_{b_\alpha^*}$. But, due to the strict concavity of the map $\eta \mapsto \log(1 - \eta)$, $\sum_{m \in \mathcal{M}} \log(1 - \eta_m^*) = \sum_{m \in \mathcal{M}} \log[1 - (\xi\eta_{1m} + (1 - \xi)\eta_{2m})] > \xi \sum_{m \in \mathcal{M}} \log(1 - \eta_{1m}) + (1 - \xi) \sum_{m \in \mathcal{M}} \log(1 - \eta_{2m}) \geq \log(1 - \alpha)$. Thus, $\eta^* \in C_\alpha \setminus UB(C_\alpha)$. By continuity of the logarithm function, there exists an $m_0 \in \mathcal{M}$ and an $\epsilon_0 > 0$ such that $\sum_{m \in \mathcal{M}} \log[1 - (\eta_m^* + \epsilon_0 I\{m = m_0\})] \geq \log(1 - \alpha)$. Observe that $(\eta_m^* + \epsilon_0 I\{m = m_0\} : m \in \mathcal{M})$ belongs to both C_α and $U(\eta^*)$. Since $\forall m \in \mathcal{M}, \eta_m \mapsto \rho_m(\eta_m)$ is strictly increasing on $C_\alpha(m)$, then $\sum_{m \in \mathcal{M}} \rho_m(\eta_m^* + \epsilon_0 I\{m = m_0\}) > \sum_{m \in \mathcal{M}} \rho_m(\eta_m^*) \geq Mb_\alpha^*$. But this contradicts the maximality of b_α^* . Therefore, $C_\alpha \cap \mathcal{N}_{b_\alpha^*}$ must be a singleton set. $\parallel$

Corollary 4.1 *If, $\forall m \in \mathcal{M}, \eta_m \in [0, \sup C_\alpha(\eta_m)) \Rightarrow \rho_m(\eta_m) < 1$, then the optimal weak FWER- α multiple decision size vector is unique.*

Proof: Follows from Theorem 4.2 and Proposition 3.1 since condition implies that for $\forall m \in \mathcal{M}, \eta_m \mapsto \rho_m(\eta_m)$ is strictly increasing on $C_\alpha(m)$. $\parallel$

Non-uniqueness of the optimal weak FWER-controlling multiple decision size vector may occur with non-regular families of densities, such as the uniform or shifted exponential densities, where the power of the test may equal one even though its size is still less than one. It may also occur if the decision processes in the MDP are not size-valid (see paragraph after Definition 3.2) such as with discrete data or nonparametric rank-based methods and randomized tests are not permitted. In such situations, the mappings $\eta_m \mapsto \rho_m(\eta_m)$ for $m \in \mathcal{M}$ need not be strictly increasing leading to a non-singleton set $C_\alpha \cap \mathcal{N}_{b_\alpha^*}$.

4.4 Finding Optimal Size Vector

This subsection addresses the computational problem of finding the optimal weak FWER multiple decision size vector. Generally, without differentiability of the ROC functions as with discrete distributions, linear or nonlinear programming methods are needed to obtain the optimal solution. Below we present a method when the ROC functions are twice-differentiable.

Theorem 4.3 *Let $\Delta^* = (\Delta_m^*, m \in \mathcal{M})$ be the most powerful MDP. Assume that the ROC functions $\eta_m \mapsto \rho_m(\eta_m)$ are strictly increasing and twice-differentiable with first and second derivatives ρ'_m and ρ''_m , respectively. Given $\alpha \in (0, 1)$, the optimal weak FWER- α multiple*

decision size vector $\eta^* \equiv \eta_{\Delta^*}^*(\alpha) = (\eta_m^*(\alpha), m \in \mathcal{M})$ is the η that solves the set of Lagrange equations

$$\forall m \in \mathcal{M}, \rho'_m(\eta_m)(1 - \eta_m) = \lambda, \text{ for some } \lambda \in \mathfrak{R}_+; \quad (4.6)$$

$$\sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha). \quad (4.7)$$

Proof: Since ρ_m s are strictly increasing, by Theorem 4.2 there is a unique optimal solution. Form the Lagrange function on $(0, 1)^M \times \mathfrak{R}$ via

$$J(\eta, \lambda) = \sum_{m \in \mathcal{M}} \rho_m(\eta_m) + \lambda \left\{ \sum_{m \in \mathcal{M}} \log(1 - \eta_m) - \log(1 - \alpha) \right\}.$$

For $m \in \mathcal{M}$, $\partial J / \partial \eta_m = \rho'_m(\eta_m) - \lambda / (1 - \eta_m)$ and $\partial J / \partial \lambda = \sum_{m=1}^M \log(1 - \eta_m) - \log(1 - \alpha)$. Equating to zeros yield conditions (4.6) and (4.7).

To show that the solution of (4.6) and (4.7) is a maximizer of J , we need to verify that the sequence of determinants of the principal minors of the bordered Hessian matrix, evaluated at this solution, alternates in signs. The second partial derivatives of the Lagrange function are, for $m, n \in \mathcal{M}$, $\partial^2 J / \partial \eta_m \partial \eta_n = \{\rho''_m(\eta_m) - \lambda / (1 - \eta_m)^2\} I\{m = n\}$, $\partial^2 J / \partial \eta_m \partial \lambda = \partial^2 J / \partial \lambda \partial \eta_m = 1 / (1 - \eta_m)$, and $\partial^2 J / \partial \lambda^2 = 0$. The solution of (4.6) and (4.7) satisfies $\lambda = \rho'_m(\eta_m)(1 - \eta_m)$, $m \in \mathcal{M}$. Since $\rho'_m(\eta_m) > 0$, then at the solution, $\lambda \geq 0$. Furthermore, since by Proposition 3.1, $\rho_m(\cdot)$ is concave, then $\rho''_m(\eta_m) \leq 0$. As a consequence, at the solution, $\partial^2 J / \partial \eta_m \partial \eta_n \leq 0$, $\partial^2 J / \partial \eta_m \partial \lambda < 0$, and $\partial^2 J / \partial \lambda^2 = 0$. The bordered $(M + 1) \times (M + 1)$ Hessian matrix evaluated at the solution of (4.6) and (4.7) is of form

$$\mathbf{H} = - \begin{bmatrix} \text{Dg}(\mathbf{b}) & \mathbf{a} \\ \mathbf{a}^t & 0 \end{bmatrix},$$

where $\text{Dg}(\mathbf{b})$ is the diagonal matrix with diagonal elements consisting of the elements of the vector $\mathbf{b}$ and with $\mathbf{b}^t = -(\rho''_m(\eta_m) - \lambda / (1 - \eta_m)^2)$, $m \in \mathcal{M}$ and $\mathbf{a}^t = (1 / (1 - \eta_m))$, $m \in \mathcal{M}$. Observe that all elements of $\mathbf{b}$ and $\mathbf{a}$ are nonnegative. The m th principal minor of $\mathbf{H}$ is

$$\mathbf{H}_m = \begin{bmatrix} \text{Dg}(\mathbf{b}_m) & \mathbf{a}_m \\ \mathbf{a}_m^t & 0 \end{bmatrix}$$

where $\mathbf{b}_m = (b_1, b_2, \dots, b_m)^t$ and $\mathbf{a}_m = (a_1, a_2, \dots, a_m)^t$. Since

$$\det(\mathbf{H}_m) = (-1)^{m+2} \left(\prod_{k=1}^m b_k \right) \sum_{k=1}^m a_k^2 / b_k$$

and the a_k s and b_k s are nonnegative, then the determinants of the principal minors of the bordered Hessian matrix alternate in sign, starting with a negative sign. Consequently, the solution of (4.6) and (4.7) is a maximizer of the Lagrange function, and hence maximizes $\sum_{m \in \mathcal{M}} \rho_m(\eta_m)$ subject to the FWER α -level constraint. $\parallel$

Proposition 4.5 *Assume the conditions of Theorem 4.3. Then, for each $m \in \mathcal{M}$, the mapping $\alpha \mapsto \eta_m^*(\alpha)$ is nondecreasing and continuous.*

Proof: Define $g_m(\eta_m) = \rho'_m(\eta_m)(1 - \eta_m)$. Since $\rho'_m(\eta_m) \geq 0$ by nondecreasing property, then $g_m(\eta_m) \geq 0, \forall \eta_m \in [0, 1]$. Furthermore, $g'_m(\eta_m) = \rho''_m(\eta_m)(1 - \eta_m) - \rho'_m(\eta_m)$, so since $\rho''_m(\eta_m) \leq 0$ by concavity, then $g'_m(\eta_m) \leq 0$. Therefore, each $g_m(\cdot)$ is a nonincreasing function. The defining condition of η^* in (4.6) is $g_m(\eta_m) = \lambda, \forall m \in \mathcal{M}$, together with the constraint condition in (4.7). Suppose α increases from α_1 to α_2 . This entails that the left-hand side of (4.7) must decrease, but for this to happen, and since each $g_m(\cdot)$ is nonincreasing, the common value of the Lagrange constant λ must decrease. But then each η_m cannot decrease. Thus, for each $m \in \mathcal{M}$, $\eta_m(\alpha_1) \leq \eta_m(\alpha_2)$. Continuity follows immediately since the existence of the second derivative implies that ρ'_m is continuous. $\parallel$

The monotonicity in Proposition 4.5 is a desirable property since it implies that if at FWER size α_1 , we have $\delta_m(\eta_m(\alpha_1)) = 1$, then at an FWER size α_2 with $\alpha_2 > \alpha_1$, we also have $\delta_m(\eta_m(\alpha_2)) = 1$. This will also be critical in proving a martingale property needed for the development of the FDR-controlling procedure. Further examination of the proofs of Propositions 4.3 and 4.4 and Theorem 4.1 also reveals that, given any MDP Δ whose ROC functions are nondecreasing, concave, and continuous, there exists an optimal FWER-controlling multiple decision size vector *specific* to this Δ . Our starting with the most powerful MDP Δ^* was in order to get the optimal FWER-controlling MDF among all MDFs in $\mathcal{D}_0$. This observation will play an important role in extending our results to situations with composite hypotheses. The basic idea is to start with a *collection* of MDPs, find the optimal weak FWER-controlling multiple decision size vector for each MDP, then choose the best among the optimal MDFs arising from each of these MDPs.

4.5 An Example for Weak FWER Control

In this subsection we demonstrate the weak FWER-controlling procedure by considering a concrete situation with normal distributions. An earlier version of the manuscript also dealt with exponential distributions and binomial distributions. However, due to space constraints, we do not present these other examples.

For $m \in \mathcal{M}$, let $X_m \sim N(\mu_m, \sigma_{m0}^2)$, where the μ_m s are unknown and σ_{m0}^2 s are known. Consider the multiple hypotheses testing problem $H_{m0} : \mu_m = \mu_{m0}$ and $H_{m1} : \mu_m = \mu_{m1}$ with $\mu_{m0} < \mu_{m1}$ for $m \in \mathcal{M}$. The MP test of size η_m for H_{m0} versus H_{m1} is

$$\delta_m^*(X_m; \eta_m) \equiv \delta_m^*(\eta_m) = I\{X_m \geq \mu_{m0} + \sigma_{m0}\Phi^{-1}(1 - \eta_m)\}, \quad (4.8)$$

where $\Phi(\cdot)$ and $\Phi^{-1}(\cdot)$ are the cumulative distribution and quantile functions, respectively, of a standard normal random variable. The m th effect size is $\gamma_m = (\mu_{m1} - \mu_{m0})/\sigma_{m0}$, and the ROC function of the decision process $\Delta_m^* = (\delta_m^*(\eta_m) : \eta_m \in [0, 1])$ is

$$\rho_m(\eta_m) \equiv \rho_m(\eta_m; \gamma_m) = \Phi(\gamma_m - \Phi^{-1}(1 - \eta_m)), \quad (4.9)$$

clearly twice-differentiable with respect to η_m . With $\phi(z) = \exp(-z^2)/\sqrt{2\pi} = \Phi'(z)$ being the standard normal density function, the derivative of $\rho_m(\cdot)$ is

$$(\rho_m)'(\eta_m) = \frac{\phi(\gamma_m - \Phi^{-1}(1 - \eta_m))}{\phi(\Phi^{-1}(1 - \eta_m))}. \quad (4.10)$$

For fixed $\alpha \in (0, 1)$ and γ_m s, consider the mappings $d \mapsto \eta_m(d)$, $m \in \mathcal{M}$, defined implicitly by the equation

$$\frac{\phi(\gamma_m - \Phi^{-1}(1 - \eta_m))}{\phi(\Phi^{-1}(1 - \eta_m))}(1 - \eta_m) - d = 0. \quad (4.11)$$

The optimal value of d , denoted by d^* , solves the equation

$$\sum_{m \in \mathcal{M}} \log(1 - \eta_m(d)) - \log(1 - \alpha) = 0. \quad (4.12)$$

The optimal sizes of the M MP tests are then $\eta_m(d^*)$, $m \in \mathcal{M}$. An R [22] implementation of this numerical problem first defines $v_m = 1 - \Phi^{-1}(1 - \eta_m)$, so condition (4.11) amounts to solving for $v_m = v_m(d)$ the equation

$$\log \Phi(v_m) + \gamma_m v_m - \log(d) - \gamma_m^2/2 = 0. \quad (4.13)$$

The R implementation utilized a Newton-Raphson iteration in solving for v_m s in (4.13) and the `uniroot` routine in the R Library to solve for d in (4.12). Upon obtaining the $v_m(d)$ s, the $\eta_m(d)$ s are computed via $\eta_m(d) = 1 - \Phi(v_m(d))$.


Figure 1 demonstrates the optimal sizes for different effect sizes when $M = 2000$ for uniformly distributed effect sizes. Observe that when the effect size is small, which converts to low power, then the optimal size for the test is also small, but also note that when the effect size is large, which converts to high power, then the optimal test size is also small. For the tests with moderate effect sizes or power, then the optimal sizes are higher. This behavior could also be seen by looking at the second panel in the figure which shows the achieved power of the tests at the optimal sizes, and in Table 1 which contains results for small values of M .

We also compared the efficiency of the optimal procedure relative to the Šidák procedure. The measure of efficiency is the ratio (multiplied by 100) of the average power over the M tests, defined by $\sum_{m \in \mathcal{M}} \rho_m(\eta_m)/M$, of the optimal procedure and the average power of the Šidák procedure. The fourth panel in Figure 1 depicts the powers of the resulting tests versus the effect size for both procedures (solid blue = optimal; dashed red = Šidák). In these uniformly-generated effect sizes, the efficiency of the optimal procedure over the Šidák is 103.5%. This efficiency is affected by the vector of effect sizes. For instance, when we change the effect sizes in Figure 1 to be generated from a uniform over $[.1, 2]$, then the efficiency jumps to 181.7%, though it should also be pointed out that since the effect sizes are small, then the overall powers of both procedures are also small.

Table 1: Optimal test sizes under normality for different power/effect size configurations. The configuration and the optimal sizes are described by the notation $k : (a, b, \dots)$ which is interpreted as having k of each of the elements in the vector $(a, b, \dots)$. The relative efficiency (in percent) of the optimal procedure relative to the Šidák procedure is also presented.

Effect Size, γ , Configuration	Optimal Test Sizes/[Efficiency over Šidák (in %)]	
	$M = 4$	$M = 20$
$M : 1$	4 : .0127 [100.0]	20 : .0026 [100.0]
$M/2 : (.5, 1)$	2 : (.0009, .0245) [113.6]	10 : (0, .0051) [125.1]
$M/2 : (1, 2)$	2 : (.0050, .0204) [104.5]	10 : (.0001, .0050) [115.3]
$M/2 : (1, 5)$	2 : (.0228, .0026) [103.6]	10 : (.0035, .0016) [100.3]
$M/4 : (0.5, 1, 2, 4)$	1 : (.0001, .0128, .0303, .0075) [105.4]	5 : (0, .0003, .0068, .0031) [107.1]
$M/4 : (1, 2, 4, 8)$	1 : (.0128, .0304, .0075, 0) [105.0]	5 : (.0003, .0068, .0031, 0) [104.3]

Figure 1: Optimal test sizes and powers for 2000 MP tests of hypotheses under normality when the effect sizes were generated from a uniform[.1, 10] distribution. Panel four shows the powers for both the optimal [solid black] and the Šidák [dashed red] tests with respect to effect sizes.



4.6 A Size-Investing Strategy

In this concrete example we observed from Figure 1 and Table 1 the phenomenon where, among the M tests, those with low powers (small effect sizes) and those with high powers (large effect sizes) are allocated relatively small sizes in the weak FWER-controlling optimal procedure. The tests getting the larger sizes are those with moderate powers or effect sizes. We refer to this as a *size-investing strategy* in the multiple hypotheses testing problem. The theoretical basis for this strategy, at least under the conditions of Theorem 4.3, is the first condition for optimality in (4.6), which is tied to the rates of change of the ROC functions of the MP multiple decision process, together with a penalty incurring from larger sizes.

This strategy can be explained intuitively. With the overall goal of getting more real discoveries while controlling the proportion of false discoveries for a pre-specified, usually small, overall size α , the optimal procedure dictates that not much size should be accorded those tests with either very low or very high powers. The former case will not lead to any discoveries anyway if the size that could be allocated is small, while the latter case will lead to discoveries even if the test sizes are made small. Thus, there is more to be gained by investing larger sizes on those tests that are of moderate powers, and an appropriate tweaking of their test sizes according to condition (4.6) improves the ability to achieve more real discoveries. However, this phenomenon is dependent on the magnitude of the overall size. If this overall size is made larger, then more leeway may ensue to the extent that it may then be more beneficial to allocate more size also to those with low powers since those tests with moderate powers, when they had small sizes, may now have larger powers because of the consequent increase in their sizes. The precise and crucial determinant of where the differential sizes should be allocated are the rates of change of the ROC functions, with some size-attenuation. See also interesting discussions of size and weight allocation strategies in [58], where the size allocation was related to the ‘ α -spending’ function of [26], and [16] which deals with α -investing in sequential procedures that control expected false discoveries, as well as [18, 34] which discuss optimal weights for the p -values.

Interestingly, a tangential real-life manifestation of this size-investing strategy occurred during the 2008 American presidential election, with the total resources (financial, manpower, etc.) available to the candidates analogous to the overall size in the multiple testing problem. In the waning days of the campaign, the major presidential candidates, then-Senator Barack Obama of the Democratic Party and Senator John McCain of the Republican Party, focussed their campaign efforts, in terms of allocating their financial and manpower resources, in the ‘battleground states’ of North Carolina, Virginia, and Pennsylvania, while basically ignoring the ‘in-the-bag states’ of South Carolina, then expected to vote for McCain, and California, then expected to vote for Obama. Also, by virtue of the deep resources of the Obama campaign, it was able to allocate more resources *even* in states that traditionally voted Republican, whereas the McCain campaign, with a relatively smaller war chest, had to ‘drop’ some states (e.g., Michigan) in their campaign. These opposing behaviors of the two camps

could be explained by the size-investing strategy with proper accounting of each campaign's overall resources.

5 Restrictions, Extensions, and Connections

5.1 On the Restriction to $\mathcal{D}_0$

The optimization problem for weak FWER control could be construed as limited since we restricted our search for the optimal MDF in the class $\mathcal{D}_0$ consisting of simple MDFs, even though the m th component of the MDF is the MP test function for H_{m0} versus H_{m1} based on (X_m, U_m) . The resulting optimal weak FWER-controlling procedure is still therefore simple. Storey [50] and Sun and Cai [53] have argued and demonstrated that performance could be improved in multiple decision problems through compound MDFs. These are characterized by the phenomenon that in testing the m th pair of hypotheses, information is borrowed from the other components of the data vector $X = (X_m : m \in \mathcal{M})$, analogous to the James-Stein [23] shrinkage phenomenon. An example of a compound MDF is the *estimated* optimal discovery procedure (ODP) in [50] and [51], though the ODP by itself as defined in Lemma 2 in [50] is not yet compound in the sense defined above, but it does use the densities of all the components to form the significance thresholding function. Other examples of compound MDFs are the FDR-controlling procedure in [1] and the oracle-based adaptive MDFs in [53].

A question arises whether we could start immediately with compound MDFs to search for an optimal weak (or strong) FWER-controlling compound MDF. Thus, suppose that $\delta = (\delta_m : m \in \mathcal{M})$ is a compound MDF so that δ_m depends on (X, U) and not only on (X_m, U_m) . For such an MDF, we have

$$R_{01}(\delta, Q) = \mathbf{P}_Q \left\{ \bigcup_{m \in \mathcal{M}_0(Q)} [\delta_m(X, U) = 1] \right\} \quad (5.1)$$

$$= 1 - \mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m(X, U) = 0] \right\}. \quad (5.2)$$

Observe that even if the independence condition (I) holds, $(\delta_m(X, U) : m \in \mathcal{M}_0(Q))$ need not be an independent collection. As such no closed-form *exact* expression for $R_{01}(\delta, Q)$ need exist. Certainly, the right-hand side expression in (5.1) could be Bonferroni-bounded by

$$\text{EFP}(\delta, Q) \equiv \sum_{m \in \mathcal{M}_0(Q)} \alpha_{\delta_m}(Q), \quad (5.3)$$

called the expected number of false positives in [50]. Alternatively, if a generalized positive quadrant dependence (PQD) condition holds among these components, which states that

$$\mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m(X, U) = 0] \right\} \geq \prod_{m \in \mathcal{M}_0(Q)} \mathbf{P}_Q \{ \delta_m(X, U) = 0 \},$$

then the right-hand side in (5.2) could be upper-bounded by

$$\text{PQD}(\delta, Q) \equiv 1 - \prod_{m \in \mathcal{M}_0(Q)} [1 - \alpha_{\delta_m}(Q)], \quad (5.4)$$

where $\alpha_{\delta_m}(Q) = E_Q \delta_m(X, U)$, the size of δ_m when $m \in \mathcal{M}_0(Q)$. For this compound MDF, its MDR is

$$R_2(\delta, Q) = \sum_{m \in \mathcal{M}_1(Q)} [1 - \pi_{\delta_m}(Q)], \quad (5.5)$$

where $\pi_{\delta_m}(Q) = E_Q \delta_m(X, U)$ is the power of δ_m when $m \in \mathcal{M}_1(Q)$.

An optimization approach is to put an upper threshold of $\alpha \in (0, 1)$ on either (5.3) or (5.4), and to obtain an MDF δ minimizing $R_2(\delta, Q)$, or, equivalently, maximizing $\text{ETP}(\delta, Q) \equiv \sum_{m \in \mathcal{M}_1(Q)} \pi_{\delta_m}(Q)$, a quantity referred in [50] as the expected number of true positives. Spjøtvoll's [46] optimal procedure is the *simple* MDF defined by

$$\delta_{SPJ}(\alpha) = \arg \max_{\delta \in \mathcal{D}_0} \{ \text{ETP}(\delta, Q_1) : \text{EFP}(\delta, Q_0) \leq \alpha \}, \quad (5.6)$$

where $Q_0 \in \mathcal{Q}_0$ and $Q_1 \in \mathcal{Q}_1$. On the other hand, Storey's [50] ODP is defined via

$$\delta_{STO}(\alpha; Q) = \arg \max_{\delta \in \mathcal{D}} \{ \text{ETP}(\delta, Q) : \text{EFP}(\delta, Q) \leq \alpha \}, \quad (5.7)$$

where Q is the true underlying probability measure of X . His use of the EFP as Type I error measure enabled a calculus of variations optimization to obtain the form of the optimal MDF. His procedure has a particularly interesting structure when we utilize as its input the vector of p -value statistics $(S_m^*(x_m, u_m) : m \in \mathcal{M})$ arising from the most powerful multiple decision process $\Delta^* = (\Delta_m^* : m \in \mathcal{M})$ with multiple decision size function $\mathbf{A}_{\Delta^*}^* = \{(A_m^*(\eta) : \eta \in [0, 1]) : m \in \mathcal{M}\}$ and multiple decision ROC function $\rho_{\Delta^*}^* = \{(\rho_m^*(\eta) : \eta \in [0, 1]) : m \in \mathcal{M}\}$ and with $A_m^*(\cdot)$ and $\rho_m^*(\cdot)$ being both differentiable with respective derivatives $(A_m^*)'(\cdot)$ and $(\rho_m^*)'(\cdot)$. Defining the so-called significance thresholding function $\mathcal{S} : ([0, 1], \sigma[0, 1]) \rightarrow \mathfrak{R}$ via

$$\mathcal{S}(s; Q) = \frac{\sum_{m \in \mathcal{M}_1(Q)} (\rho_m^*)'(s)}{\sum_{m \in \mathcal{M}_0(Q)} (A_m^*)'(s)}, \quad (5.8)$$

with this expression following directly from Lemma 2 in [50] and Proposition 3.2, then $\delta_{STO} = (\delta_{m, STO} : m \in \mathcal{M})$ has the structure

$$\delta_{m, STO}(S_m^*(x_m, u_m); Q) = I\{\mathcal{S}(S_m^*(x_m, u_m); Q) \geq \lambda\}, \quad m \in \mathcal{M}, \quad (5.9)$$

for $\lambda \in [0, \infty)$ chosen so the size constraint on $\text{EFP}(\delta_{STO}(\alpha; Q), Q)$ is *approximately* satisfied. Observe that each of the components in (5.9) is *still* of simple-type, that is, the m th component depends only on (x_m, u_m) , unless the cut-off λ is determined in a data-dependent

manner using the full data (x, u) . Note also that δ_{STO} was derived under complete knowledge of the unknown Q , or more specifically, the sets $\mathcal{M}_0(Q)$ and $\mathcal{M}_1(Q)$, as can be seen in (5.8). For the simple null versus simple alternative case, the size functions $A_m^*(\cdot)$ s and the ROC functions $\rho_m^*(\cdot)$ s will be known, but with composite hypothesis they may be unknown. In order to implement δ_{STO} , it was proposed in [50] and [51] that these unknown quantities, sets, functions, or the significance thresholding function, be estimated based on the data (x, u) . But, by doing so, which now makes the MDF of compound-type, the exact optimality property of the ODP need not hold anymore, though it could be argued that if good estimates are utilized then the resulting estimated ODP will have desirable properties. See [53] and [15] for an interesting discussion of the ODP procedure. In contrast, note that δ_{SPJ} is determined only by the extreme probability measures Q_0 and Q_1 whose marginal probability measures, Q_m s, are completely known, and not by the unknown true probability measure Q . This fact was criticized by Storey [50] as a ‘potentially problematic optimality’ criterion. More importantly, it should be recognized that both δ_{SPJ} and δ_{STO} are not necessarily the optimal weak or strong FWER- or FDR-controlling MDFs since the Bonferroni upper bound for $R_{01}(\delta, Q)$ utilized in the derivations is hardly a sharp upper bound.

The criticism leveled against δ_{SPJ} could also be invoked against our optimal weak FWER-controlling procedure since in our optimization we also relied on a criterion that was determined only by the extreme probability measures Q_0 and Q_1 . Note, however, that *each* of the components of the optimal weak FWER-controlling multiple decision size vector, and consequently each of the components of $\delta_W^*(\alpha)$, uses *all* of the Q_{m0} s and Q_{m1} s, analogously to Storey’s ODP, though the MDF $\delta_W^*(\alpha)$ is at this point neither adaptive nor compound. Our development of this simple MDF, which is optimal in the class $\mathcal{D}_0$, is a prelude to the development of our adaptive and compound MDFs *strongly*-controlling FWER and FDR. The MDF $\delta_W^*(\alpha)$ serves as anchor towards the development of these FWER and FDR strongly-controlling compound MDFs. These new MDFs will be discussed in Section 6 for the strong FWER-control and in Section 7 for the strong FDR control. One may characterize our approach to obtaining these strongly-controlling MDFs as indirect, whereas Storey’s [50] approach maybe viewed as a more direct approach. There is also an intrinsic difference in the problems considered since we are focussing on the Type I error risk functions R_{01} and R_1 , whereas in [50] and [46] the simpler Type I error metric of expected number of false positives (EFP) was utilized. Looking forward, though our starting point is still the optimal weak FWER-controlling simple MDF $\delta_W^*(\alpha)$, there is confidence in the viability of this indirect approach to generate good MDFs since, as will be demonstrated later, the sequential Šidák procedure and the BH procedure are special cases of the strong FWER- and FDR-controlling compound MDFs, both arising under exchangeability.

5.2 Families with MLR Property

The initial simplification of the multiple decision problem to the simple null versus simple alternative hypotheses for each $m \in \mathcal{M}$ may be perceived as a serious limitation since the optimal MDF will then depend, and hence requires knowledge, of the Q_{m1} s to calculate the ROC functions. In settings with “large M , small n ” data sets, such knowledge may not be so forthcoming. However, we point out that the simplistic approach of simply assuming, most probably erroneously, that the (Q_{m0}, Q_{m1}) is invariant in $m \in \mathcal{M}$, which is the *exchangeable setting*, will have undesirable consequences; see [13]. Historically, in the development of optimal procedures, such as in the Neyman-Pearson hypothesis testing framework, it is prudent to start with the simplest setting, which usually turns out to be the most fundamental setting. In our situation this is the case with simple null and simple alternative hypotheses. This approach was also implemented in [46], [50], and [34]. Recall that in the development of optimal classes of test functions in the single-pair hypothesis testing problem, the role of the MP test is centrally crucial. We surmise that in the multiple decision problem, the solution to the simple null versus simple alternative hypotheses problem will play a prominent role in solving the composite hypotheses setting. It appears that for an MTP to possess optimality, it will require knowledge, either exact, approximate, or estimated, of the alternative hypotheses distributions; see also [34]. We briefly touch on this aspect in the presence of the monotone likelihood ratio (MLR) property, see [28].

Consider the situation where, for each $m \in \mathcal{M}$, the density function q_m belongs to a one-dimensional parametric family $\mathcal{F}_m = \{q_m(\cdot; \xi_m) : \xi_m \in \Gamma_m \subset \mathbb{R}\}$ which possesses the MLR property. A typical pair of hypotheses to be tested would be $H_{m0}^* : \xi_m \leq \xi_{m0}$ versus $H_{m1}^* : \xi_m > \xi_{m0}$, where ξ_{m0} is known. With the MLR property, a uniformly most powerful (UMP) test function $\delta_m(X_m, U_m; \eta_m)$ of size η_m exists, with this UMP test identical to the MP test of size η_m for the simple null hypothesis $H_{m0} : \xi_m = \xi_{m0}$ versus the simple alternative hypothesis $H_{m1} : \xi_m = \xi_{m1}$, with $\xi_{m1} > \xi_{m0}$. When dealing with the single-pair hypothesis testing problem, we recall that exact knowledge of the value of ξ_1 is not necessary since the critical constants of the size- η MP test for $H_0 : \xi = \xi_0$ versus $H_1 : \xi = \xi_1$ can be made independent of ξ_1 . In contrast, for the multiple decision problem, to determine the optimal size allocations for each of the M MP tests, for a given overall weak FWER threshold, the powers of the tests at the ξ_{m1} s are required, hence the need to know the values of the ξ_{m1} s. We propose two possible solutions to this dilemma.

The first approach is to solicit from the scientific investigator the values of the ξ_{m1} s for which the powers are of most interest. Such values may coincide with those that are scientifically, e.g., clinically, different from the ξ_{m0} s. Such elicitation, which may not be very feasible in practice if M is large, but which may be made possible by forming subclasses or clusters of the M genes as in [13], amounts to specifying *effect sizes*, analogous to that in sample size determination problems. Formation of such clusters must be made in close consultation with the investigator, or perhaps guided by the result of a preliminary cluster

analysis using data independent of that used in the decision functions. For the specified ξ_{m1} s, the ROC functions in the determination of the optimal weak FWER-controlling multiple size vector become $\rho_m(\eta) = \pi_{\delta_m^*(\eta)}(\xi_{m1})$ for $m \in \mathcal{M}$, where $\delta_m^*(\eta)$ is the simple MP test of size η for testing $H_{m0} : \xi_m = \xi_{m0}$ versus $H_{m1} : \xi_m = \xi_{m1}$, and $\pi_{\delta_m^*(\eta)}(\xi_{m1})$ is the power of $\delta_m^*(\eta)$ (at $\xi_m = \xi_{m1}$). In the clustered situation with $\mathcal{M} = \biguplus_{k=1}^K \mathcal{M}_k$, where $(\mathcal{M}_k, k = 1, 2, \dots, K)$ partitions $\mathcal{M}$, we may denote by $\bar{\rho}_k(\eta)$ and ζ_k , respectively, the common ROC function and size for the decision functions in cluster $\mathcal{M}_k$. Under second-order differentiability of $\bar{\rho}_k(\eta)$ s, by Theorem 4.3, the optimal weak FWER- α controlling multiple size vector $\zeta(\alpha) = (\zeta_1(\alpha), \zeta_2(\alpha), \dots, \zeta_K(\alpha))$ is the $\zeta = (\zeta_1, \zeta_2, \dots, \zeta_K)$ that solves the set of equations

$$\begin{aligned} \forall k = 1, 2, \dots, K : \bar{\rho}'_k(\zeta_k)(1 - \zeta_k) &= \lambda \text{ for some } \lambda \in \mathbb{R}; \\ \sum_{k=1}^K |\mathcal{M}_k| \log(1 - \zeta_k) &= \log(1 - \alpha). \end{aligned}$$

The second approach, which is analogous to what has been done in several papers such as [58], [36], [53], [50], [51], and [25], is to estimate or approximate the underlying values of the ξ_m s either using the observed data x , possibly using shrinkage-type estimators, or through the use of prior information, possibly informed by external covariates as in [15]. Addressing this same restriction of requiring knowledge of the simple null and simple alternative hypotheses and advocating this second approach, Roquain and van de Wiel [34] wrote: “Although leading to oracle procedures, it can be used in practice as soon as the null and alternative distributions are estimated or guessed reasonably accurately from independent data.” By ‘independent data’ is meant in [34] as data different from that used in performing the actual tests. However, such *external* data need not always be used for estimating or imputing the unknown parameters or distributions. For example, let us suppose that for each $m \in \mathcal{M}$, data x_m could be partitioned into (v_m, w_m) . We may then use $\hat{\xi}_m(v_m) = \max\{\xi_{m0}, \hat{\xi}_m(v_m)\}$, where $\hat{\xi}_m(v_m)$ is the maximum likelihood estimate of ξ_m based on v_m . We may then proceed as in the preceding paragraph where ξ_{m1} is set to $\hat{\xi}_m(v_m)$ for each $m \in \mathcal{M}$, but with the component data w_m used in the test functions. The resulting MDF will then be of an adaptive type, possibly also compound, similar to those in [53], if shrinkage estimators are used for estimating the ξ_m s using the v_m components. Observe that if for some $m_0 \in \mathcal{M}$, $\hat{\xi}_{m_0}(v_{m_0})$ and ξ_{m_0} are very close or identical, then a relatively small size will be allocated to the MP test for component m_0 . (Refer to subsection 4.6 for a discussion on the behavior of the components of the optimal multiple decision size vector.) This then amounts to downgrading, or simply ignoring, the testing problem for this component, a fact that is of importance since a criticism of multiple hypotheses testing, especially when using the FDR error rate, is that an unscrupulous investigator may just keep adding irrelevant genes. When using the adaptive MDF arising from the optimal multiple decision size vector, this investigator’s strategy will be foiled since the adaptive MDF will automatically downgrade the irrelevant genes.

This second approach, however, still requires further study. For instance, there is the issue of how to partition each x_m into the v_m and w_m components. Furthermore, the impact

of a misspecified ξ_{m1} , possibly arising from the estimation procedure, needs to be ascertained. For example, suppose that the true ξ_{m1} is approximated by $\tilde{\xi}_{m1}$. Then, the optimal weak FWER-controlling multiple size vector will be robust to such misspecification provided that the mappings, for $m = 1, 2, \dots, M$,

$$\eta_m \mapsto \frac{\rho_m^{11}(\eta_m; \xi_{m1})}{\rho_m^{10}(\eta_m; \xi_{m1})}(\tilde{\xi}_{m1} - \xi_{m1})$$

are approximately invariant in m . Here, $\rho_m(\eta; \xi_{m1})$ is the ROC function associated with the MP decision process for testing $H_{m0} : \xi_m = \xi_{m0}$ versus $H_{m1} : \xi_m = \xi_{m1}$, and $\rho_m^{ab}(\eta; \xi) = \{\partial^{a+b}/\partial\eta^a\partial\xi^b\}\rho_m(\eta; \xi)$. However, we are still currently studying these and related issues pertaining to the implementation of the procedures in more depth.

An alternative procedure, but which we do not advocate, is to utilize the full data x to estimate the ξ_m s, and then to utilize again x in the decision functions. However, this approach could have unacceptable consequences due to re-using or double-dipping on the data x . Such re-use of the data may invalidate desired properties of the MDF, such as control of the Type I error rate; see the discussion in [15].

5.3 Connections to p -Value Statistics

Corollary 3.2 indicates that the ROC function $\eta \mapsto \rho_m(\eta)$ is differentiable if and only if the distribution function of the, possibly randomized, p -value statistic $S_m(X_m, U_m)$ under $H_{m1} : Q_m = Q_{m1}$ is differentiable. In this case, $\rho'_m(\cdot)$ coincides with $h_m(\cdot)$, the density function of $S_m(X_m, U_m)$ under $H_{m1} : Q_m = Q_{m1}$. The first condition (4.6) in Theorem 4.3 may then be restated in terms of these density functions via

$$h_m(\eta_m)(1 - \eta_m) = \text{Constant}, \quad \forall m \in \mathcal{M}. \quad (5.10)$$

This is a surprising result as it indicates that it is *not* enough to simply find the sizes that maximize these $h_m(\cdot)$ s, as dictated by the Neyman-Pearson Lemma when dealing with a single pair of null and alternative hypotheses. Rather, in the multiple hypotheses testing problem, there is attenuation in that larger sizes incur some penalties. This is the phenomenon referred to in subsection 4.6 as a size-investing strategy. Equation (5.10) governs the interactions among the M tests regarding their size allocations to achieve the best overall result, in terms of overall Type II error, among themselves. See also the last paragraph in section 3 of [34] regarding the form of their optimal multi-weighted step-up procedure.

The optimal weak FWER-controlling MDF may be converted to a procedure based on the p -value statistics. If $\eta^*(\alpha) = (\eta_m^*(\alpha), m \in \mathcal{M})$ is the optimal weak FWER- α multiple decision size vector and $(S_m(x_m, u_m), m \in \mathcal{M})$ is the vector of computed p -value statistics, the decision based on data $(x, u) = ((x_m, u_m), m \in \mathcal{M})$ is $\delta^*(x, u) = (I\{S_m(x_m, u_m) \leq \eta_m^*(\alpha)\}, m \in \mathcal{M})$, an MDF based on weighted p -values. This is related to the approach

proposed in several papers of using weighted p -values such as in [18], [55], [36], [25], and [34]. In our case the weights are tied-in to the optimal sizes. Observe that it is possible to have $S_{m_1}(x_{m_1}, u_{m_1}) < S_{m_2}(x_{m_2}, u_{m_2})$ but with $H_{m_1 0}$ not rejected and $H_{m_2 0}$ rejected depending on the values of $\eta_{m_1}^*(\alpha)$ and $\eta_{m_2}^*(\alpha)$. In such cases, decision-making for the M pairs of hypotheses need not be transitive with respect to the p -value statistics; see section 3 in [53] where this behavior also occurs with their oracle-based adaptive procedure for control of mFDR.

6 Strong FWER Control

In this section we develop a strong FWER-controlling compound MDF using as a starting point the optimal weak FWER-controlling procedure developed in Section 4.

Let $\Delta^* = (\Delta_m^*, m \in \mathcal{M})$ be the MP MDP with $\Delta_m^* = (\delta_m^*(\eta) : \eta \in [0, 1])$ the MP decision process for $H_{m0} : Q_m = Q_{m0}$ versus $H_{m1} : Q_m = Q_{m1}$ based on (X_m, U_m) . Without loss of generality, we assume that the size function $A_m(\cdot)$ of Δ_m^* satisfies $A_m(\eta) = \eta$. Define the mapping $\eta : [0, 1] \rightarrow [0, 1]^M$ such that $\eta(\alpha) = (\eta_m(\alpha), m \in \mathcal{M})$ is the optimal weak FWER-controlling multiple decision size vector at level α . We shall assume in this section that each component of this mapping is nondecreasing and continuous, which is the case for instance when the ROC functions of Δ^* are twice-differentiable as established in Proposition 4.5.

For an FWER threshold of $\alpha \in [0, 1]$, the optimal MDF in $\mathcal{D}_0$ is

$$\delta_W^*(\alpha) = (\delta_m^*(\eta_m(\alpha)), m \in \mathcal{M}). \quad (6.1)$$

Associated with this MDF is the *generalized* multiple decision p -value statistic $\mathbf{W} = (W_m, m \in \mathcal{M})$, where

$$W_m \equiv W_m(X_m, U_m) = \inf\{\alpha \in [0, 1] : \delta_m^*(\eta_m(\alpha)) = 1\}. \quad (6.2)$$

The $w_m = W_m(x_m, u_m)$ is the smallest weak FWER size leading to rejection of H_{m0} when using $\delta_W^*(\alpha)$ given data $(x, u) = ((x_m, u_m), m \in \mathcal{M})$. The usual p -value statistic S_m [see (3.3)] for δ_m^* is related to W_m via

$$\forall m \in \mathcal{M} : S_m(X_m, U_m) = \eta_m(W_m(X_m, U_m)). \quad (6.3)$$

In the sequel, the statistic $\mathbf{Q}$, which takes values in the set of permutations of $(1, 2, \dots, M)$, and denoted by

$$\mathbf{Q} = (Q_1, Q_2, \dots, Q_M) = ((1), (2), \dots, (M)), \quad (6.4)$$

represents the anti-rank vector of $\mathbf{W}$, so that $W_{Q_1} < W_{Q_2} < \dots < W_{Q_M}$ or, equivalently, $W_{(1)} < W_{(2)} < \dots < W_{(M)}$.

Now, a la [50, 53], suppose an Oracle knows Q , the true underlying probability measure governing X . For the MDF $\delta_W^*(\alpha)$ in (6.1), its FWER is

$$R_{01}(\delta_W^*(\alpha), Q) = 1 - \prod_{m \in \mathcal{M}} [1 - \eta_m(\alpha)]^{1 - \theta_m(Q)}.$$

This is a nondecreasing and continuous function of α by virtue of the assumed nondecreasingness and continuity properties of the mappings $\alpha \mapsto \eta_m(\alpha)$ for each $m \in \mathcal{M}$. Now, if the Oracle desires to control this Type I error rate at a value $q^* \in [0, 1]$ and also minimize the MDR given by

$$R_2(\delta_W^*(\alpha), Q) = |\mathcal{M}_1(Q)| - \sum_{m \in \mathcal{M}_1(Q)} \rho_m(\eta_m(\alpha))$$

where $\rho_m(\eta_m(\alpha))$ is the power of $\delta_m^*(\eta_m(\alpha))$, then She should choose the largest $\alpha \in [0, 1]$ such that $R_{01}(\delta_W^*(\alpha), Q) = q^*$. Owing to the continuity and nondecreasingness of $R_{01}(\delta_W^*(\alpha), Q)$ in α , the Oracle's optimal α could also be expressed via

$$\alpha^\dagger(q^*; Q) = \inf \left\{ \alpha \in [0, 1] : \prod_{m \in \mathcal{M}} [1 - \eta_m(\alpha)]^{1 - \theta_m(Q)} < 1 - q^* \right\},$$

with the convention that $\inf \emptyset = 1$.

However, there is no Oracle and Q is not known, else there is no multiple decision problem. So $\alpha^\dagger(q^*; Q)$ is not observable! A natural idea is to estimate the unknown $\theta_m(Q)$, the state of the m th pair of hypotheses. An intuitive and simple estimator of $\theta_m(Q)$ for a fixed value of α is

$$\hat{\theta}_m(Q) = \delta_m^*(\eta_m(\alpha)-) \equiv \delta_m^*(X_m, U_m; \eta_m(\alpha)-). \quad (6.5)$$

In turn we obtain a *step-down* estimator $\alpha^\dagger(q^*) \equiv \alpha^\dagger(X, U; q^*)$ of the Oracle-based $\alpha^\dagger(q^*; Q)$ given by

$$\alpha^\dagger(q^*) = \inf \left\{ \alpha \in [0, 1] : \prod_{m \in \mathcal{M}} [1 - \eta_m(\alpha)]^{1 - \delta_m^*(\eta_m(\alpha)-)} < 1 - q^* \right\}. \quad (6.6)$$

This then determines the compound MDF $\delta_S^*(q^*) \equiv \delta_S^*(X, U; q^*) \in \mathcal{D}$, where

$$\delta_S^*(q^*) = (\delta_m^*(\eta_m(\alpha^\dagger(q^*))), m \in \mathcal{M}). \quad (6.7)$$

By virtue of the optimal choice of the $\eta_m(\alpha)$ s and the use of the MP tests, we expect $\delta_S^*(q^*)$ to possess excellent, if not optimal, MDR-properties. By taking the infimum over the weak FWER-size α coupled with the estimation of $\theta_m(Q)$ by $\delta_m^*(\eta_m(\alpha)-)$ in (6.6), there occurs an adaptive downweighting of components whose H_{m0} s are most likely correct as dictated by the data (x, u) . We now establish that $\delta_S^*(q^*)$ in (6.7) do control strongly the FWER.

Theorem 6.1 *Let $q^* \in [0, 1]$. Then, $\forall Q \in \mathcal{Q}$, $R_{01}(\delta_S^*(q^*), Q) \leq q^*$.*

Proof: Fix a $q^* \in [0, 1]$ and let $Q \in \mathcal{Q}$ be the true underlying probability measure of X . Define the stochastic process $\mathbf{T}_1 = \{T_1(\alpha) : \alpha \in [0, 1]\}$ via

$$T_1(\alpha) = \prod_{m \in \mathcal{M}} [1 - \eta_m(\alpha)]^{1 - \delta_m^*(\eta_m(\alpha)-)}.$$

The sample paths of $\mathbf{T}_1$ are, a.e. $[Q]$, left-continuous with right-hand limits (caglad), piecewise nonincreasing, and with $T_1(\alpha-) = T_1(\alpha) \leq T_1(\alpha+)$. In terms of $\mathbf{T}_1$, we have $\alpha^\dagger(q^*) = \inf\{\alpha \in [0, 1] : T_1(\alpha) < 1 - q^*\}$. Consequently, $T_1(\alpha^\dagger(q^*)) \geq 1 - q^*$. Now, note that

$$R_{01}(\delta_S^*(q^*), Q) = 1 - \mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m^*(\eta_m(\alpha^\dagger(q^*))) = 0] \right\}. \quad (6.8)$$

Observe that we could not write the last term as a product since the events concerned are not anymore independent owing to the fact that $\alpha^\dagger(q^*)$ depends on the whole data (X, U) . We also have that

$$\bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m^*(\eta_m(\alpha^\dagger(q^*))) = 0] = \left\{ \alpha^\dagger(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \right\}.$$

Next, define the stochastic process $\mathbf{T}_2 = \{T_2(\alpha) : \alpha \in [0, 1]\}$ via

$$T_2(\alpha) = \left(\prod_{m \in \mathcal{M}_0(Q)} [1 - \eta_m(\alpha)] \right) \left(\prod_{m \in \mathcal{M}_1(Q)} [1 - \eta_m(\alpha)]^{1 - \delta_m^*(\eta_m(\alpha)-)} \right).$$

This process, which depends on the unknown Q , also has caglad sample paths. Let

$$\alpha^\#(q^*) = \alpha^\#(q^*; Q) = \inf\{\alpha \in [0, 1] : T_2(\alpha; Q) < 1 - q^*\}.$$

In contrast to $\alpha^\dagger(q^*)$, $\alpha^\#(q^*)$ is not a random variable since it depends on the unknown Q . Note that we also have $T_2(\alpha^\#(q^*)) \geq 1 - q^*$. It is also clear from their definitions that $T_1(\alpha) \geq T_2(\alpha; Q)$. As such, since $T_1(\alpha) < 1 - q^*$ implies $T_2(\alpha; Q) < 1 - q^*$, then it follows that $\alpha^\dagger(q^*) \geq \alpha^\#(q^*)$. More importantly, observe that

$$\left\{ \alpha^\dagger(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \right\} = \left\{ \alpha^\#(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \right\}.$$

To see this, the inclusion $\subseteq$ is immediate from the last result. Since $\{\alpha^\#(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m\}$ implies that for some $\alpha_0 < \min_{m \in \mathcal{M}_0(Q)} W_m$, then we have $T_2(\alpha_0; Q) < 1 - q^*$. For such an α_0 , $\delta_m(\eta_m(\alpha_0)-) = 0$ for all $m \in \mathcal{M}_0(Q)$. Thus, $\alpha_0 \in \{\alpha \in [0, 1] : T_1(\alpha) < 1 - q^*\}$. Consequently, $\alpha^\dagger(q^*) \equiv \inf\{\alpha \in [0, 1] : T_1(\alpha) < 1 - q^*\} \leq \alpha_0 < \min_{m \in \mathcal{M}_0(Q)} W_m$. Thus, $\supseteq$ inclusion holds, completing the proof of the set equality.

Invoking the iterated expectation rule and the above equivalences, we have

$$\begin{aligned}
& \mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m^*(\eta_m(\alpha^\dagger(q^*))) = 0] \right\} \\
&= \mathbf{P}_Q \left\{ \alpha^\dagger(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \right\} \\
&= \mathbf{P}_Q \left\{ \alpha^\#(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \right\} \\
&= E_Q \left[\mathbf{P}_Q \left\{ \alpha^\#(q^*) < \min_{m \in \mathcal{M}_0(Q)} W_m \mid \alpha^\#(q^*) \right\} \right].
\end{aligned}$$

The quantity $\alpha^\#(q^*)$ is measurable with respect to $\sigma\{\delta_m^* : m \in \mathcal{M}_1(Q)\}$, whereas the random variable $\min_{m \in \mathcal{M}_0(Q)} W_m$ is measurable with respect to $\sigma\{\delta_m^* : m \in \mathcal{M}_0(Q)\}$. These sub-sigma-fields are independent since δ_m^* s are simple decision functions and by virtue of the independence condition (I). As a consequence,

$$\begin{aligned}
\mathbf{P}_Q \left\{ \min_{m \in \mathcal{M}_0(Q)} W_m > w \right\} &= \mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m^*(\eta_m(w)) = 0] \right\} \\
&= \prod_{m \in \mathcal{M}_0(Q)} \mathbf{P}_Q \left\{ \delta_m^*(\eta_m(w)) = 0 \right\} = \prod_{m \in \mathcal{M}_0(Q)} [1 - \eta_m(w)],
\end{aligned}$$

with the product arising because of the independence of the $\{\delta_m^* : m \in \mathcal{M}_0(Q)\}$ from condition (I) and since the δ_m^* s are simple. Therefore,

$$\begin{aligned}
& \mathbf{P}_Q \left\{ \bigcap_{m \in \mathcal{M}_0(Q)} [\delta_m^*(\eta_m(\alpha^\dagger(q^*))) = 0] \right\} \\
&= E_Q \left\{ \prod_{m \in \mathcal{M}_0(Q)} [1 - \eta_m(\alpha^\#(q^*))] \right\} \\
&\geq E_Q \{T_2(\alpha^\#(q^*))\} \\
&\geq E_Q(1 - q^*) \\
&= 1 - q^*.
\end{aligned}$$

Using this result and (6.8), we have $R_{01}(\delta_S^*(q^*), Q) \leq 1 - (1 - q^*) = q^*$, proving the theorem. $\parallel$

Let us relate $\delta_S^*(q^*)$ to the p -value and generalized p -value statistics $\mathbf{S}$ and $\mathbf{W}$, respectively. Define the random variable

$$\begin{aligned}
J^\dagger(q^*) &= \max \{j \in \mathcal{M} : T_1(W_{(i)}) \geq 1 - q^*, i = 1, 2, \dots, j\} \\
&= \max \left\{ j \in \mathcal{M} : \prod_{m=i}^M [1 - \eta_{(m)}(W_{(i)})] \geq 1 - q^*, i = 1, 2, \dots, j \right\}.
\end{aligned}$$

Then, $\alpha^\dagger(q^*) \in [W_{(J^\dagger(q^*))}, W_{(J^\dagger(q^*)+1)}]$, so we may re-express $\delta_S^*(q^*)$ via

$$\delta_S^*(q^*) = (\delta_m^*(\eta_m(W_{(J^\dagger(q^*))})), m \in \mathcal{M}).$$

Suppose now that we are in the exchangeable setting where $Q_{m0} = Q_0$ and $Q_{m1} = Q_1$ for all $m \in \mathcal{M}$, so that the ROC functions for the M decision processes are identical. The optimal weak FWER-controlling multiple decision size vector will have identical components and will coincide with the Šidák sizes, that is, for each $m \in \mathcal{M}$, $\eta_m(\alpha) = 1 - (1 - \alpha)^{1/M} = \eta_m^S(\alpha)$. From (6.3) we then have that for each $m \in \mathcal{M}$,

$$S_{(m)} = 1 - (1 - W_{(m)})^{1/M} \quad \text{and} \quad W_{(m)} = 1 - (1 - S_{(m)})^M.$$

It follows that

$$\begin{aligned} J^\dagger(q^*) &= \max \left\{ j \in \mathcal{M} : \prod_{m=i}^M (1 - W_{(i)})^{1/M} \geq 1 - q^*, i = 1, 2, \dots, j \right\} \\ &= \max \left\{ j \in \mathcal{M} : (1 - W_{(i)})^{(M-i+1)/M} \geq 1 - q^*, i = 1, 2, \dots, j \right\} \\ &= \max \left\{ j \in \mathcal{M} : W_{(i)} \leq 1 - (1 - q^*)^{M/(M-i+1)}, i = 1, 2, \dots, j \right\} \\ &= \max \left\{ j \in \mathcal{M} : S_{(i)} \leq 1 - (1 - q^*)^{1/(M-i+1)}, i = 1, 2, \dots, j \right\}. \end{aligned}$$

Therefore, in terms of the usual p -value statistics $(S_1, S_2, \dots, S_M)$, H_{m0} will be rejected if and only if the rank of S_m is at most $J^\dagger(q^*)$. This is precisely the FWER-controlling step-down Šidák compound MDF; see, for instance, Procedure 3.9 on page 123 in Dudoit and van der Laan [9]. We have thus established that the sequential step-down Šidák procedure is a special case of the compound MDF $\delta_S^*(q^*)$, arising under the exchangeable setting.

7 Strong FDR Control

In this section we obtain an FDR-controlling compound MDF anchored on the weak FWER-controlling simple MDF $\delta_W^*(\alpha)$ in (6.1). We consider the same framework as in Section 6. In particular we assume that, for each $m \in \mathcal{M}$, the mapping $\alpha \in [0, 1] \mapsto \eta_m(\alpha)$ is nondecreasing and continuous, where $\eta(\alpha) = (\eta_m(\alpha), m \in \mathcal{M})$ is the optimal weak FWER- α multiple size vector. Our idea in obtaining an FDR-controlling MDF builds on the development of the BH MDF, specifically Benjamini and Hochberg's [1] rationale of their Theorem 2.

Let $q^* \in [0, 1]$ be the desired FDR threshold and Q be the underlying probability measure of X . We introduce two stochastic processes: $\mathbf{T}_0 = \{T_0(\alpha; Q) : \alpha \in [0, 1]\}$ and $\mathbf{T} = \{T(\alpha) : \alpha \in [0, 1]\}$, where

$$T_0(\alpha; Q) = \sum_{m \in \mathcal{M}_0(Q)} \delta_m^*(\eta_m(\alpha)) \quad \text{and} \quad T(\alpha) = \sum_{m \in \mathcal{M}} \delta_m^*(\eta_m(\alpha)).$$

For the MDF $\delta_W^*(\alpha)$, its FDR is

$$R_1(\delta_W^*(\alpha), Q) = E_Q \left\{ \frac{T_0(\alpha; Q)}{T(\alpha)} I\{T(\alpha) > 0\} \right\}.$$

By definition of the generalized p -value statistics W_m s in (6.2) we have for $\alpha \in [W_{(m)}, W_{(m+1)})$ that $T(\alpha) = m$, whereas

$$E_Q\{T_0(\alpha; Q)\} = \sum_{m \in \mathcal{M}} (1 - \theta_m(Q)) \eta_m(\alpha) \leq \sum_{m \in \mathcal{M}} \eta_m(\alpha). \quad (7.1)$$

Suppose we focus on an $\alpha \in [W_{(m)}, W_{(m+1)})$. If $\sum_{j \in \mathcal{M}} \eta_j(W_{(m)}) \leq mq^*$, then the best α in this interval will be the largest value satisfying $\sum_{j \in \mathcal{M}} \eta_j(\alpha) \leq mq^*$, since by increasing α , the MDR decreases as argued in the development of $\delta_S^*(q^*)$ in Section 6. This motivates our definition of $\alpha^*(q^*) = \alpha^*(X, U; q^*)$ as the *step-up* estimator

$$\alpha^*(q^*) = \sup \left\{ \alpha \in [0, 1] : \sum_{m \in \mathcal{M}} \eta_m(\alpha) \leq q^* \sum_{m \in \mathcal{M}} \delta_m^*(\eta_m(\alpha)) \right\}. \quad (7.2)$$

This induces a compound MDF $\delta_F^*(q^*) \equiv \delta_F^*(X, U; q^*) \in \mathcal{D}$ given by

$$\delta_F^*(q^*) = (\delta_m^*(\eta_m(\alpha^*(q^*))), m \in \mathcal{M}). \quad (7.3)$$

Another possibility, which may lead to further improvements, is to estimate $\theta_m(Q)$ in (7.1) by $\delta_m^*(\eta_m(\alpha)-)$, as in Section 6, and replace $\sum_{m \in \mathcal{M}} \eta_m(\alpha)$ in (7.2) by $\sum_{m \in \mathcal{M}} [1 - \delta_m^*(\eta_m(\alpha)-)] \eta_m(\alpha)$. This may extend the adaptive procedure in [2]; however, we defer its consideration for future work. We establish that $\delta_F^*(q^*)$ controls the FDR at q^* .

Theorem 7.1 *Let $q^* \in [0, 1]$. If, $\forall Q \in \mathcal{Q} \setminus \{Q_0\}$ and $\forall \alpha \in (0, 1)$,*

$$|\mathcal{M}_0(Q)| \max_{m \in \mathcal{M}_0(Q)} \eta_m(\alpha) \leq \sum_{m \in \mathcal{M}} \eta_m(\alpha), \quad (7.4)$$

then $R_1(\delta_F^(q^*), Q) \leq q^*$ for $\forall Q \in \mathcal{Q}$.*

Proof: The cases $q^* = 0$ and $q^* = 1$ are clearly trivial, so let $q^* \in (0, 1)$, and let $Q \in \mathcal{Q}$ be the underlying probability measure of X . First, we note that for $\forall \alpha \in [0, 1]$, $\eta_m(\alpha) \leq \alpha$ for all $m \in \mathcal{M}$. For each $\alpha \in (0, 1]$, let us define the sigma-field

$$\mathcal{F}_\alpha = \sigma\{\delta_m^*(X_m, U_m; \eta_m(\beta)) : m \in \mathcal{M}, \alpha \leq \beta \leq 1\},$$

and let $\mathcal{F}_0 = \bigvee_{\alpha \in (0, 1]} \mathcal{F}_\alpha$. Observe that for $0 \leq \alpha < \beta \leq 1$, $\mathcal{F}_\alpha \supset \mathcal{F}_\beta$. Denote by $\mathbf{F} = \{\mathcal{F}_\alpha : \alpha \in [0, 1]\}$ the induced filtration. For conciseness, we shall drop (X_m, U_m) in $\delta_m^*(X_m, U_m; \eta_m)$

and simply write $\delta_m^*(\eta_m)$, and also write α^* for $\alpha^*(q^*)$. Observe that the stochastic processes $\mathbf{T}_0$ and $\mathbf{T}$ defined earlier are $\mathbf{F}$ -adapted.

For the MDF $\delta_F^* \equiv \delta_F^*(q^*)$ in (7.3), its FDR is

$$R_1(\delta_F^*, Q) = E_Q \left[E_Q \left\{ \frac{T_0(\alpha^*; Q)}{T(\alpha^*)} I\{T(\alpha^*) > 0\} | \mathcal{M}_0 \right\} \right],$$

where, given Q , $\mathcal{M}_0 = \mathcal{M}_0(Q)$ is a random, albeit degenerate, subset of $\mathcal{M}$. Thus, the outer expectation is an expectation with respect to this degenerate probability measure. Let us focus on the inner expectation. If $M_0 = |\mathcal{M}_0| = 0$, then this expectation is zero since $T_0(\alpha^*) = 0$, hence bounded by q^* .

Consider the case where $M_0 \in \{1, 2, \dots, M-1\}$. From the definition of $\alpha^* = \alpha^*(q^*)$ in (7.2), we see that it is a $\mathbf{F}$ -stopping time. Also, observe that at $\alpha = \alpha^*$, we have the inequality $T(\alpha^*) \geq \sum_{m \in \mathcal{M}} \eta_m(\alpha^*)/q^* \equiv \eta_\bullet(\alpha^*)/q^*$ with $\eta_\bullet(\alpha) = \sum_{m \in \mathcal{M}} \eta_m(\alpha)$. Consequently,

$$\begin{aligned} E_Q \left\{ \frac{T_0(\alpha^*)}{T(\alpha^*)} I\{T(\alpha^*) > 0\} | \mathcal{M}_0 \right\} &\leq q^* E_Q \left\{ \frac{T_0(\alpha^*)}{\eta_\bullet(\alpha^*)} I\{T(\alpha^*) > 0\} | \mathcal{M}_0 \right\} \\ &= q^* E_Q \left\{ \frac{T_0(\alpha^*)}{\eta_\bullet(\alpha^*)} | \mathcal{M}_0 \right\}, \end{aligned} \tag{7.5}$$

the last equality following since $T_0(\alpha) I\{T(\alpha) > 0\} = T_0(\alpha)$ for every $\alpha \in [0, 1]$. Next, consider the $\mathbf{F}$ -adapted process $\mathbf{T}_0^* = \{T_0^*(\alpha) : \alpha \in (0, 1]\}$ with

$$T_0^*(\alpha) \equiv T_0^*(\alpha; Q) = \sum_{m \in \mathcal{M}_0(Q)} \frac{\delta_m^*(\eta_m(\alpha))}{\eta_m(\alpha)}.$$

We have, for $0 < \alpha < \beta \leq 1$, that

$$\begin{aligned}
E_Q\{T_0^*(\alpha)|\mathcal{F}_\beta, \mathcal{M}_0\} &= E_Q\left\{\sum_{m \in \mathcal{M}_0} \frac{\delta_m^*(\eta_m(\alpha))}{\eta_m(\alpha)}|\mathcal{F}_\beta, \mathcal{M}_0\right\} \\
&= \sum_{m \in \mathcal{M}_0} E_Q\left\{\frac{\delta_m^*(\eta_m(\alpha))}{\eta_m(\alpha)}|\mathcal{F}_\beta, \mathcal{M}_0\right\} \\
&= \sum_{m \in \mathcal{M}_0} \delta_m^*(\eta_m(\beta)) E_Q\left\{\frac{\delta_m^*(\eta_m(\alpha))}{\eta_m(\alpha)}|\delta_m^*(\eta_m(\beta)) = 1, \mathcal{F}_\beta, \mathcal{M}_0\right\} \\
&= \sum_{m \in \mathcal{M}_0} \delta_m^*(\eta_m(\beta)) E_{Q_{m0}}\left\{\frac{\delta_m^*(\eta_m(\alpha))}{\eta_m(\alpha)}|\delta_m^*(\eta_m(\beta)) = 1\right\} \\
&= \sum_{m \in \mathcal{M}_0} \frac{\delta_m^*(\eta_m(\beta))}{\eta_m(\alpha)} \frac{\mathbf{P}_{Q_{m0}}\{\delta_m^*(\eta_m(\alpha)) = 1\}}{\mathbf{P}_{Q_{m0}}\{\delta_m^*(\eta_m(\beta)) = 1\}} \\
&= \sum_{m \in \mathcal{M}_0} \frac{\delta_m^*(\eta_m(\beta))}{\eta_m(\alpha)} \frac{\eta_m(\alpha)}{\eta_m(\beta)} \\
&= \sum_{m \in \mathcal{M}_0} \frac{\delta_m^*(\eta_m(\beta))}{\eta_m(\beta)} \\
&= T_0^*(\beta),
\end{aligned}$$

where the fourth equality follows from Condition (I) in (2.3). The fifth equality follows from the assumed nondecreasing property of the mapping $\alpha \mapsto \eta_m(\alpha)$, which implies that for $\alpha < \beta$, $\{\delta_m^*(\eta_m(\alpha)) = 1\} \Rightarrow \{\delta_m^*(\eta_m(\beta)) = 1\}$, and the size condition $\mathbf{P}_{Q_{m0}}\{\delta_m^*(\eta_m) = 1\} = E_{Q_{m0}}[\delta_m^*(\eta_m)] = \eta_m$ on the MP tests or the decision processes Δ_m^* s. This sequence of equalities establishes the result that $\{(T_0^*(\alpha), \mathcal{F}_\alpha) : \alpha \in (0, 1]\}$ is a reverse martingale process. Define $T_0^*(0) = \liminf_{\alpha \in (0, 1]} T_0^*(\alpha) = \limsup_{\alpha \in (0, 1]} T_0^*(\alpha)$, which, by Doob's martingale convergence theorem, is well-defined. Then the extended collection $\{(T_0^*(\alpha), \mathcal{F}_\alpha) : \alpha \in [0, 1]\}$ is

a reverse martingale process. The expectation portion of the upper bound in (7.5) becomes

$$\begin{aligned}
E_Q \left\{ \frac{T_0(\alpha^*)}{\eta_{\bullet}(\alpha^*)} | \mathcal{M}_0 \right\} &= E_Q \left\{ \frac{\sum_{m \in \mathcal{M}_0} \delta_m^*(\eta_m(\alpha^*))}{\eta_{\bullet}(\alpha^*)} | \mathcal{M}_0 \right\} \\
&= E_Q \left\{ \sum_{m \in \mathcal{M}_0} \frac{\delta_m^*(\eta_m(\alpha^*))}{\eta_m(\alpha^*)} \frac{\eta_m(\alpha^*)}{\eta_{\bullet}(\alpha^*)} | \mathcal{M}_0 \right\} \\
&\leq \left[\sup_{\alpha \in (0,1]} \frac{\max_{m \in \mathcal{M}_0} \eta_m(\alpha)}{\eta_{\bullet}(\alpha)} \right] E_Q \{ T_0^*(\alpha^*) | \mathcal{M}_0 \} \\
&\leq \frac{1}{M_0} E_Q \{ T_0^*(\alpha^*) | \mathcal{M}_0 \} \\
&= \frac{1}{M_0} E_Q \{ T_0^*(1) | \mathcal{M}_0 \} \\
&= \frac{1}{M_0} \sum_{m \in \mathcal{M}_0} \frac{E_{Q_{m0}} \{ \delta_m^*(\eta_m(1)) | \mathcal{M}_0 \}}{\eta_m(1)} \\
&= \frac{1}{M_0} \sum_{m \in \mathcal{M}_0} \frac{\eta_m(1)}{\eta_m(1)} \\
&= 1,
\end{aligned}$$

where the condition (7.4) was used to get the second inequality and the Optional Sampling Theorem for martingales [6] to get the third equality. Thus, we have established that

$$E_Q \left\{ \frac{T_0(\alpha^*)}{T(\alpha^*)} I \{ T(\alpha^*) > 0 \} | \mathcal{M}_0 \right\} \leq q^*$$

for any $\mathcal{M}_0$ provided $|\mathcal{M}_0| = M_0 \leq M - 1$. Taking expectation with respect to $\mathcal{M}_0$, which as pointed out earlier is a degenerate probability measure, yields the result that $R_1(\delta_F^*, Q) \leq q^*$.

Finally, consider $M_0 = M$, that is, all H_{m0} s are correct. Recall the Šidák size vector for weak FWER control at α given by

$$\eta_m^S(\alpha) = \eta_1^S(\alpha) = 1 - (1 - \alpha)^{1/M}, \quad m \in \mathcal{M}.$$

The vector $\eta^S(\alpha) = (\eta_m^S(\alpha) : m \in \mathcal{M})$ clearly satisfies condition (7.4). With $\eta_{\bullet}^S(\alpha) = M\eta_1^S(\alpha)$, let $\alpha^S \equiv \alpha^S(q^*) = \alpha^S(X, U; q^*)$ be

$$\alpha^S = \sup \left\{ \alpha \in [0, 1] : \eta_{\bullet}^S(\alpha) \leq q^* \sum_{m \in \mathcal{M}} \delta_m^*(\eta_m^S(\alpha)) \right\} \quad (7.6)$$

and its associated MDF given by $\delta^S = (\delta_m^*(\eta_m^S(\alpha)), m \in \mathcal{M})$. Since in the proof for the case with $M_0 \in \{0, 1, 2, \dots, M - 1\}$ it is not necessary that the size vector $(\eta_m(\alpha), m \in \mathcal{M})$ be

the weak FWER-controlling optimal size vector, then the proof also holds when we use the Šidák size vector. Furthermore, since the Šidák size vector satisfies condition (7.4) even when $M_0 = M$, then for the Šidák sizes, we have for all $Q \in \mathcal{Q}$, including Q_0 , that $R_1(\delta^S, Q) \leq q^*$. Define, for $\alpha \in [0, 1]$, the processes

$$V^*(\alpha) = \frac{\sum_{m \in \mathcal{M}} \delta_m^*(\eta_m(\alpha))}{\sum_{m \in \mathcal{M}} \eta_m(\alpha)} \quad \text{and} \quad V^S(\alpha) = \frac{\sum_{m \in \mathcal{M}} \delta_m^*(\eta_m^S(\alpha))}{\sum_{m \in \mathcal{M}} \eta_m^S(\alpha)}$$

with $V^*(0) = V^S(0) = 0$. Observe that, under Q_0 , the expectations of $V^*(\alpha)$ and $V^S(\alpha)$ are both equal to 1 for each $\alpha \in (0, 1]$. We may re-express both α^* and α^S via

$$\begin{aligned} \alpha^* &= \sup\{\alpha \in [0, 1] : V^*(\alpha) \geq 1/q^*\}; \\ \alpha^S &= \sup\{\alpha \in [0, 1] : V^S(\alpha) \geq 1/q^*\}. \end{aligned}$$

Since $q^* < 1$, by Lemma 7.1, which is stated and established below, it follows that

$$\mathbf{P}_{Q_0} \left\{ V^*(\alpha) \geq \frac{1}{q^*} \right\} \leq \mathbf{P}_{Q_0} \left\{ V^S(\alpha) \geq \frac{1}{q^*} \right\}.$$

This implies that, under Q_0 , $\alpha^* \stackrel{st}{\leq} \alpha^S$. From this it follows that

$$R_1(\delta_F^*, Q_0) = \mathbf{P}_{Q_0}(\alpha^* > 0) \leq P_{Q_0}(\alpha^S > 0) = R_1(\delta^S, Q_0) \leq q^*.$$

Though not essential in the proof, notice that, in fact, $R_1(\delta^S, Q_0) = q^*$. This is seen by noting that under the Šidák size vector η^S and the associated MDF δ^S , all the inequalities in the proof are in fact equalities. We have thus completed the proof that, whatever Q is, $R_1(\delta^*, Q) \leq q^*$. $\parallel$

Lemma 7.1 *Let $(V_m, m \in \mathcal{M})$ be independent random variables with $V_m \sim \text{Ber}(\eta_m)$ where $\eta = (\eta_m, m \in \mathcal{M}) \in [0, 1]^M$. For $a \geq 0$, define*

$$h_a(\eta) = \mathbf{P} \left\{ \frac{\sum_{m=1}^M V_m}{\sum_{m=1}^M \eta_m} \geq a \right\}.$$

Then, $\forall \alpha \in (0, 1)$ and $\forall a \in [1, \infty)$, $\sup \{h_a(\eta) : \eta \in UB(C_\alpha)\} = h_a(\eta^S(\alpha))$.

Proof: Let $Z_1, Z_2, \dots, Z_M$ be independent random variables with $Z_m \sim \text{Ber}(p_m)$ and denote by $\bar{p} = \frac{1}{M} \sum_{m=1}^M p_m$. For $t \geq 0$, let

$$h_t^*(p_1, p_2, \dots, p_M) = \mathbf{P} \left\{ \sum_{m=1}^M Z_m \geq t \right\}.$$

In Hoeffding [21] (see also pages 375–376 of Marshall and Olkin [30]) it was proved that if $M\bar{p} \leq t \leq M$, then $h_t^*(p_1, p_2, \dots, p_M) \leq h_t^*(\bar{p}, \bar{p}, \dots, \bar{p})$. In the setting of the lemma, define $p_m = p_m(\eta) = -\log(1-\eta_m)$ for $m \in \mathcal{M}$. Then, $\eta \in UB(C_\alpha)$ iff $\sum_{m=1}^M p_m(\eta_m) = -\log(1-\alpha)$. For $a \geq 1$, we are then able to apply the result in [21] to conclude that for all $\eta \in UB(C_\alpha)$,

$$\begin{aligned} h_a(\eta) &= h_{-\frac{a}{M}\log(1-\alpha)}^*(p_1(\eta_1), \dots, p_m(\eta_m)) \\ &\leq h_{-\frac{a}{M}\log(1-\alpha)}^*\left(-\frac{1}{M}\log(1-\alpha), \dots, -\frac{1}{M}\log(1-\alpha)\right) = h_a(\eta^S). \end{aligned}$$

This proves the lemma. $\parallel$

We establish that $\delta_F^*(q^*)$ reduces to the BH procedure under the exchangeable setting.

Corollary 7.1 *If the ROC functions are identical, then $\delta_F^*(q^*)$ is the FDR- q^* controlling MDF in Benjamini and Hochberg [1].*

Proof: The condition implies that $\forall m \in \mathcal{M}, \eta_m(\alpha) = \eta(\alpha)$ for some $\eta(\alpha)$. Therefore, (7.2) becomes

$$\alpha^*(q^*) = \sup \left\{ \alpha \in [0, 1] : M\eta(\alpha) \leq q^* \sum_{m \in \mathcal{M}} \delta_m^*(\eta(\alpha)) \right\}. \quad (7.7)$$

Relabeling $\eta(\alpha)$ by just α in (7.7) and noting that each $\delta_m^*(\alpha)$ could be re-expressed via $\delta_m^*(\alpha) = I\{S_m(X_m, U_m) \leq \alpha\}$, then (7.7) becomes

$$\alpha^S(q^*) = \sup \left\{ \alpha \in [0, 1] : \alpha \leq \frac{q^*}{M} \sum_{m=1}^M I\{S_m(X_m, U_m) \leq \alpha\} \right\}.$$

With $S_{(1)} < S_{(2)} < \dots < S_{(M)}$ denoting the ordered p -value statistics, we may define the random variable, as in [1],

$$J^S(q^*) = J^S(X, U; q^*) = \max \left\{ m \in \mathcal{M}^* : S_{(m)} \leq \frac{q^* m}{M} \right\}. \quad (7.8)$$

It is then easy to see that $\alpha^S(q^*) \in [S_{(J^S(q^*))}, S_{(J^S(q^*)+1)})$. Now, when using the MDF $\delta_F^*(q^*)$, H_{m0} is rejected if and only if $S_m \leq \alpha^S(q^*)$, but from the above relation, this occurs if and only if the rank of S_m is no more than $J^S(q^*)$. This latter procedure is precisely the BH FDR- q^* controlling MDF in [1]. $\parallel$

Notice in the preceding proof that the BH MDF, denoted $\delta^{BH}(q^*)$, coincides with the Šidák-size based MDF $\delta^S(q^*)$. The martingale proof for Theorem 7.1 thus carries over to establishing strong FDR control by $\delta^{BH}(q^*)$. A martingale-based proof of FDR control by $\delta^{BH}(q^*)$ was also in [52]. Furthermore, we point out that the MDF $\delta_F^*(q^*)$ fulfills, in an

exact manner, the FDR constraint. In contrast, the estimated ODP MDF in [50] only *approximately* satisfies this constraint.

Some remarks are in order regarding condition (7.4). As noted above, the Šidák multiple decision size vector, which could be viewed as the optimal multiple decision size vector when the ROC functions are identical, always satisfies (7.4). In general, when not in the exchangeable setting, condition (7.4) induces a form of control of the differences of these ROC functions. We conjecture that a weaker condition is possible to still achieve FDR-control by the MDF δ_F^* . But a non-martingale-based proof may be needed to resolve this. We also provide an alternative form of $\delta_F^*(q^*)$ in terms of the generalized p -value statistics W_m s. Define

$$J^*(q^*) \equiv J^*(X, U; q^*) = \max \left\{ m \in \mathcal{M} : \sum_{j \in \mathcal{M}} \eta_j(W_{(m)}) \leq q^* m \right\}. \quad (7.9)$$

Then $\delta_F^*(q^*)$ rejects $H_{(m)0}$ for $m \in \{1, 2, \dots, J^*(q^*)\}$ and accepts $H_{(m)0}$ for $m \in \{J^*(q^*) + 1, J^*(q^*) + 2, \dots, M\}$.

Let us also examine further the W_m s. Focussing on $W_{(1)}$, under Q_0 , we have, for $a \in (0, 1)$, $\mathbf{P}_{Q_0}(W_{(1)} > a) = \mathbf{P}_{Q_0} \left\{ \bigcap_{m \in \mathcal{M}} [\delta_m^*(\eta_m(a)) = 0] \right\} = \prod_{m \in \mathcal{M}} [1 - \eta_m(a)] = 1 - a$, using the independence of the δ_m^* s under Q_0 . Thus, $W_{(1)}$ is standard uniform when all null hypotheses are correct. Using this uniformity result and the following lemma about lower and upper bounds of $\eta_\bullet$ for $\eta \in UB(C_\alpha)$, we obtain in Proposition 7.1 a lower bound for $R_1(\delta_F^*(q^*), Q_0)$, which is the FDR when all the null hypotheses are correct.

Lemma 7.2 *Every $\eta \in UB(C_\alpha)$ satisfies*

$$\alpha \leq \eta_\bullet = \sum_{m \in \mathcal{M}} \eta_m \leq \min \left\{ -\log(1 - \alpha), M[1 - (1 - \alpha)^{1/M}] \right\}.$$

Proof: For $\eta \in UB(C_\alpha)$, let $V_1, V_2, \dots, V_M$ be independent Bernoulli random variables with $V_m \sim \text{Ber}(\eta_m)$. Bonferroni's inequality yields

$$\begin{aligned} \alpha &= 1 - \prod_{m \in \mathcal{M}} [1 - \eta_m] = \mathbf{P} \left\{ \bigcup_{m \in \mathcal{M}} [V_m = 1] \right\} \\ &\leq \sum_{m \in \mathcal{M}} \mathbf{P}\{V_m = 1\} = \sum_{m \in \mathcal{M}} \eta_m = \eta_\bullet, \end{aligned}$$

establishing the left-hand inequality.

Since for every $a \in [0, 1)$, $-\log(1 - a) \geq a$, then from the constraint condition $\sum_{m \in \mathcal{M}} \log(1 - \eta_m) = \log(1 - \alpha)$, we obtain $\eta_\bullet \leq -\log(1 - \alpha)$. But since $a \mapsto \log(1 - a)$ is concave in $[0, 1)$, then

$$\log\{(1 - \alpha)^{1/M}\} = \frac{1}{M} \sum_{m \in \mathcal{M}} \log(1 - \eta_m) \leq \log \left(1 - \frac{1}{M} \sum_{m \in \mathcal{M}} \eta_m \right).$$

This implies that $(1 - \alpha)^{1/M} \leq 1 - \eta_{\bullet}/M$ which is equivalent to $\eta_{\bullet} \leq M[1 - (1 - \alpha)^{1/M}]$. $\parallel$

Proposition 7.1 $\forall q^* \in [0, 1], 1 - (1 - q^*/M)^M \leq R_1(\delta_F^*(q^*), Q_0) \leq q^*$.

Proof: It remains to show the left-hand inequality. We have that

$$\begin{aligned} R_1(\delta_F^*(q^*), Q_0) &= \mathbf{P}_{Q_0}(\alpha^*(q^*) > 0) \\ &= \mathbf{P}_{Q_0} \left\{ \bigcup_{m=1}^M [\eta_{\bullet}(W_{(m)}) \leq mq^*] \right\} \geq \mathbf{P}_{Q_0} \{\eta_{\bullet}(W_{(1)}) \leq q^*\}. \end{aligned}$$

Using the first component of the upper bound for $\eta_{\bullet}$ in Lemma 7.2, we get $\mathbf{P}_{Q_0} \{\eta_{\bullet}(W_{(1)}) \leq q^*\} \geq \mathbf{P}_{Q_0} \{-\log(1 - W_{(1)}) \leq q^*\} = 1 - \exp(-q^*)$ since $-\log(1 - W_{(1)})$ is unit exponential owing to the standard uniformity of $W_{(1)}$ under Q_0 . Using the second component in the upper bound, we also get

$$\begin{aligned} \mathbf{P}_{Q_0} \{\eta_{\bullet}(W_{(1)}) \leq q^*\} &\geq \mathbf{P}_{Q_0} \{M[1 - (1 - W_{(1)})^{1/M}] \leq q^*\} \\ &= \mathbf{P}_{Q_0} \left\{ \alpha_{(1)} \leq 1 - (1 - q^*/M)^M \right\} = 1 - (1 - q^*/M)^M. \end{aligned}$$

Thus, $R_1(\delta_F^*(q^*), Q_0) \geq \max\{1 - \exp(-q^*), 1 - (1 - q^*/M)^M\}$. But, as a referee had pointed out, or which we see by noting that $v \in [0, \infty) \mapsto L(v) \equiv M \log(1 - v/M) + v$ has $L(0) = 0$ and $L'(v) < 0$ for $v > 0$, we have for $q^* \in [0, \infty)$, $1 - \exp(-q^*) \leq 1 - (1 - q^*/M)^M$. Thus, this completes the proof of the proposition. $\parallel$

8 A Modest Simulation

In this section we present results of a modest simulation comparing the performance of the δ_F^* and δ^{BH} in terms of FDR and MDR. The results here are limited to demonstrating numerically, in a specific Gaussian model, that δ_F^* achieves the desired FDR-control, as does δ^{BH} , and that δ_F^* achieves a lower MDR relative to δ^{BH} .

The simulation model is similar to the Gaussian example illustrating the optimal weak FWER-controlling procedure. In this model, for each $m \in \mathcal{M}$, the observables are $X_m \sim N(\mu_m, 1)$ which are independently generated. The m th pair of hypotheses is $H_{m0} : \mu_m \leq 0$ versus $H_{m1} : \mu_m > 0$ with UMP size- η_m test of form $\delta_m^*(X_m; \eta_m) = I\{X_m > \Phi^{-1}(1 - \eta_m)\}$. The true values of the means μ_m s are $\mu_m = \xi_m \theta_m, m \in \mathcal{M}$, with $\theta_m \sim \text{Ber}(p)$ and effect sizes $\xi_m \sim |N(\nu, 1)|$, which are again independently generated from each other. In the simulation, the parameter combinations were induced by taking the number of pairs of hypotheses $M \in \{20, 50, 100\}$, the proportion of true alternative hypotheses $p \in \{.1, .2, .4\}$, and the mean of the effect size-generating normal distribution $\nu \in \{1, 2, 4\}$. In implementing δ_F^* and δ^{BH} , we used an FDR-threshold of $q^* \in \{.05, .10\}$. Since the computational implementation of

δ_F^* takes time, for each combination of (q^*, M, ν, p) , we only replicated the basic experiment 1000 times. For each simulation parameter combination, the simulated FDR and MDR* were the averages of the observed FDR and the standardized $\text{MDR}^* = \text{MDR}/|\mathcal{M}_1(Q)|$ over the 1000 replications. For summarization purposes, we used this standardized MDR since, for each replicate, a Q is generated, hence $|\mathcal{M}_1(Q)|$ may differ over the simulation replications. Thus, in essence, note that we are comparing the averages of $R_2(\delta_F^*, Q)/|\mathcal{M}_1(Q)|$ and $R_2(\delta^{BH}, Q)/|\mathcal{M}_1(Q)|$, where the averaging is with respect to the mechanism generating the Q s over the simulation replications.

We only report the results for $q^* = 0.10$ in Table 2 since the results for $q^* = 0.05$ lead to similar conclusions. From this table we observe that both δ_F^* and δ^{BH} fulfill the FDR-constraint, and in fact this happens in a conservative fashion, which is as expected from theory. More importantly, the MDR-performance of δ_F^* is better compared to that of δ^{BH} , and this dominance holds true for all the twenty-seven simulation parameter combinations considered. Observe that as M is increased with (ν, p) remaining the same, there is an increase in their MDR^* s; whereas, when ν is increased, which has the effect of increasing the effect sizes, their MDR^* s decrease. Interestingly, the impact of a change of value in p , the proportion of true alternative hypotheses, did not necessarily translate into a monotone change in their MDR^* s, especially when $M = 20$, though for the larger M -values, the change in MDR^* appears monotonically decreasing.

It may appear that the standardized improvement of δ_F^* over δ^{BH} is minuscule based on the results of this simulation study. However, it should be noted that when translated to the overall number of discoveries, when M is large, δ_F^* will lead to many more discoveries than δ^{BH} while still maintaining the desired FDR control. Such an increase in the number of discoveries may have important practical implications, such as enlarging the number of genes that will be explored in consequent studies, hence increasing the chances of finding crucial and important genes, but without sacrificing the Type I error rate.

9 Concluding Remarks

This paper provides some resolution on the role of the individual powers, or more appropriately the ROC functions of decision processes, in multiple hypotheses testing problems. The importance and relevance of these problems is evident as witnessed by the explosion in the number of research papers that were published, and certainly those that were not published, on this subject in the last few years. A primary impetus for this development is the urgent need to deal with the proliferation of high-dimensional “large M , small n ” data sets in the natural, medical, physical, economic, and social sciences, which are being created or generated due to advances in high-throughput technology, the latter fueled by speedy developments in computer technology and miniaturization. This is embodied and spearheaded by, but not limited to, microarray technology.

Table 2: Comparison of the false discovery rate (FDR) and standardized missed discovery rate (MDR*) performance of MHTDF δ^* and δ^{BH} under a variety of simulation parameters. This table is for $q^* = .10$. The FDR and MDR* are in percentages. The number of replications is 1000.

	q^*	M	ν	p	δ^* -FDR	δ^* -MDR*	δ^{BH} -FDR	δ^{BH} -MDR*
1	0.1	20	1	0.1	8.03	70.80	8.43	72.64
2	0.1	20	1	0.2	7.55	79.64	8.77	81.99
3	0.1	20	1	0.4	6.05	77.47	6.65	80.30
4	0.1	20	2	0.1	7.70	54.42	8.43	55.80
5	0.1	20	2	0.2	7.39	56.32	7.59	57.31
6	0.1	20	2	0.4	6.47	47.82	6.21	49.38
7	0.1	20	4	0.1	9.14	8.62	9.48	10.30
8	0.1	20	4	0.2	7.80	7.34	6.97	9.20
9	0.1	20	4	0.4	6.15	3.58	5.65	5.53
10	0.1	50	1	0.1	8.83	84.87	9.26	87.05
11	0.1	50	1	0.2	7.11	83.49	7.14	86.65
12	0.1	50	1	0.4	6.45	78.91	6.42	82.30
13	0.1	50	2	0.1	8.36	63.36	8.99	65.04
14	0.1	50	2	0.2	8.74	57.30	8.73	58.93
15	0.1	50	2	0.4	5.80	48.71	5.93	50.21
16	0.1	50	4	0.1	8.84	10.28	8.93	12.09
17	0.1	50	4	0.2	7.93	6.91	7.81	8.79
18	0.1	50	4	0.4	6.34	3.40	6.07	5.68
19	0.1	100	1	0.1	9.14	87.10	9.02	90.02
20	0.1	100	1	0.2	8.21	84.05	8.78	87.38
21	0.1	100	1	0.4	5.92	80.12	5.88	83.73
22	0.1	100	2	0.1	9.79	66.10	9.24	67.93
23	0.1	100	2	0.2	7.68	58.25	7.94	59.93
24	0.1	100	2	0.4	5.74	49.29	6.10	50.90
25	0.1	100	4	0.1	8.37	10.44	8.62	12.36
26	0.1	100	4	0.2	7.72	5.93	7.81	8.22
27	0.1	100	4	0.4	5.69	3.80	6.14	5.72

Almost a century ago, Neyman and Pearson demonstrated the need to take into account the power function, and the alternative hypothesis configuration, when one is seeking an optimal test procedure in the one-pair hypothesis testing problem. Their work led to a divorce from the then-existing significance or p -value approach. Currently, many multiple hypotheses testing procedures, epitomized by the Šidák procedures for weak and strong control of the FWER, and by the well-known Benjamini-Hochberg (BH) procedure for control of the FDR, are based on the p -values of the individual tests and do not seem to consider possible differences in the powers of the individual tests. They are appropriate in the so-called exchangeable setting wherein powers of the individual tests are identical.

In this paper we examined the question of whether differences in power characteristics of the individual tests could be exploited to improve on existing procedures for FWER and FDR control. This was done in a general decision-theoretic framework to allow for results that are applicable even with complicated data types and structures, and in the most fundamental setting where each pair of hypotheses consists of a simple null and a simple alternative hypothesis. First, an optimal MDF within the class of simple MDFs was shown to exist for weak FWER control. This MDF exploits differences in the power characteristics of the individual tests. In particular, this MDF is better than the Šidák weak FWER-controlling MDF, though the latter is a special case of the optimal MDF arising under the exchangeable setting. The resulting theory also informs us regarding an optimal size-investing strategy. Second, by using this optimal, though still restricted, MDF as an anchor, we developed a compound MDF which strongly controls the FWER. The sequential Šidák MDF is a special case of this MDF, arising under the exchangeable setting. We then developed a compound MDF that (strongly) controls the FDR. The BH FDR-controlling MDF is a special case, arising under exchangeability. These new MDFs, by virtue of their construction, are expected to have smaller MDRs compared to those which do not exploit power differences. This was demonstrated through a modest simulation study for the new FDR-controlling MDF. The MDFs were also related and contrasted with other compound MDFs, notably the ODP in [50], and those using weighted p -values.

Though the proposed MDFs do improve on existing ones developed under the exchangeable setting, we could not claim that they are optimal among *all* compound MDFs for control of FWER or FDR. This question of global optimality appears to be a difficult and elusive problem. So far none of the existing compound MDFs, such as the *estimated* ODP in [50], could claim global optimality. In our case, the possible drawback is the fact that in the construction of these new MDFs, the starting point is the class of simple MDFs. Indeed, the resulting MDFs are compound, but the issue of establishing global optimality is not transparent. In fact, a question even arise as to whether there actually exists an optimal MDF among all compound MDFs that, say, control the FDR. One thing certain about our proposed MDFs is that they do satisfy the desired FWER or FDR constraints. Other MDFs, obtained by plugging-in estimators hence are adaptive, or which utilize prior information, may lose their optimality property or may not anymore satisfy desired Type I error con-

straints after the plug-in step. See [53] where optimality was in an asymptotic sense and with the Type I error rate being the mFDR, as well as [15] and [34] for more discussions on these issues.

A natural layer to add in the decision-theoretic formulation of the problem is a Bayesian layer where a prior measure is specified on the unknown probability measure Q or, alternatively, on $\theta(Q)$. There is a possibility that through this Bayesian approach, one may be able to obtain a characterization of the class of optimal MDFs controlling Type I error rates, or when the two types of error rates are combined, for example, via a weighted linear combination. The papers [31, 39, 12, 13] which employ Bayes or empirical Bayes approaches are certainly highly relevant.

Finally, we mention that there are still other aspects of the multiple decision problem not dealt with in this paper. The first one is the extension to situations with composite null and alternative hypotheses, or how to adapt the new MDFs to such settings. We indicated some ideas in subsection 5.2 for distributional models possessing the MLR property, but clearly more extensive studies are needed. The second one is that of possible dependencies among the components of $(X_m, m \in \mathcal{M}_0(Q))$. In the setting we considered, it was assumed that this is an independent collection according to condition (I), but it would be of interest to obtain results under certain types of dependencies. Potential results in such scenarios will extend those in [37, 38] and [3]. In both of these settings, resampling-based ideas and approaches, such as the use of permutational distributions, which were developed and implemented, for instance, in [57] and [56] will be highly relevant.

Acknowledgements

Many ideas and results were obtained while the first author was a Faculty Fellow at the Statistical and Applied Mathematical Sciences Institute (SAMSI) during his sabbatical leave from the University of South Carolina last Fall 2008. He is grateful to SAMSI Director Dr. James Berger for facilitating his stay at the Institute. As such, this work was partially supported by the National Science Foundation (NSF) under Grant DMS-0635449 to SAMSI. However, any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. Other results were obtained when the first author visited the University of Oslo and the Norwegian University of Science and Technology (NTNU). He wishes to thank Professor Odd Aalen and Professor Bo Lindqvist for facilitating his visit. We are grateful to the two reviewers, the Associate Editor, and the Editor for their comments, suggestions, and criticisms which increased our understanding and appreciation of multiple decision problems and led to a better manuscript. Special thanks to Prof. Sanat Sarkar for detecting a problem with our initial version of a strong FWER-controlling procedure, a problem also pinpointed by a referee, and for his many comments regarding the manuscript and for helpful discus-

sions. We also thank Prof. Lan Wang for her careful reading of the manuscript, and thank also the following colleagues for comments and/or pointing out relevant references: Prof. B. Lindqvist, Prof. J. Lynch, Dr. A. McLain, Prof. G. Rempala, Prof. J. Sethuraman, Prof. G. Taraldsen, Prof. A. Vidyashankar, Prof. L. Wasserman, and Prof. P. Westfall. Nevertheless, any errors in the manuscript are entirely by the authors. Finally, we thank Dr. M. Peña for discussions about microarray technology.

References

- [1] BENJAMINI, Y. AND HOCHBERG, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B* **57**, 1, 289–300.
- [2] BENJAMINI, Y. AND HOCHBERG, Y. (2000). On the adaptive control of the false discovery rate in multiple testing with independent statistics. *Journal of Educational and Behavioral Statistics* *25*, 60–83.
- [3] BENJAMINI, Y. AND YEKUTIELI, D. (2001). The control of the false discovery rate in multiple testing under dependency. *Ann. Statist.* **29**, 4, 1165–1188.
- [4] BONFERRONI, C. (1937). Teoria statistica delle classi e calcolo delle probabilita. In *Volume in Onore di Ricarrdo dalla Volta*. Universita di Firenze, 1–62.
- [5] COX, D. R. AND HINKLEY, D. V. (1974). *Theoretical statistics*. Chapman and Hall, London.
- [6] DOOB, J. L. (1953). *Stochastic processes*. John Wiley & Sons Inc., New York.
- [7] DUDOIT, S., GILBERT, H. N., AND VAN DER LAAN, M. (2007). Resampling-based empirical bayes multiple testing procedures for controlling generalized tail probability and expected value error rates: Focus on the false discovery rate and simulation study. Tech. rep., University of California at Berkeley.
- [8] DUDOIT, S., SHAFFER, J. P., AND BOLDRICK, J. C. (2003). Multiple hypothesis testing in microarray experiments. *Statist. Sci.* **18**, 1, 71–103.
- [9] DUDOIT, S. AND VAN DER LAAN, M. J. (2008). *Multiple testing procedures with applications to genomics*. Springer Series in Statistics. Springer, New York.
- [10] EFRON, B. (2004). Large-scale simultaneous hypothesis testing: the choice of a null hypothesis. *Journal of the American Statistical Association* *99*, 96–104.
- [11] EFRON, B. (2007). Size, power and false discovery rates. *Ann. Statist.* **35**, 4, 1351–1377.
- [12] EFRON, B. (2008a). Microarrays, empirical Bayes and the two-groups model. *Statist. Sci.* **23**, 1, 1–22.

- [13] EFRON, B. (2008b). Simultaneous inference: When should hypothesis testing problems be combined? *The Annals of Applied Statistics* **1**, 197–223.
- [14] EFRON, B., TIBSHIRANI, R., STOREY, J. D., AND TUSHER, V. (2001). Empirical Bayes analysis of a microarray experiment. *J. Amer. Statist. Assoc.* **96**, 456, 1151–1160.
- [15] FERKINGSTAD, E., FRIGESSI, A., RUE, H., THORLEIFSSON, G., AND KONG, A. (2008). Unsupervised empirical bayesian multiple testing with external covariates. *The Annals of Applied Statistics* **2**, 714–735.
- [16] FOSTER, D. P. AND STINE, R. A. (2008). α -investing: a procedure for sequential control of expected false discoveries. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **70**, 2, 429–444. <http://dx.doi.org/10.1111/j.1467-9868.2007.00643.x>.
- [17] GENOVESE, C. AND WASSERMAN, L. (2002). Operating characteristic and extensions of the false discovery rate procedure. *Journal of the Royal Statistical Society, Series B* **64**, 499–517.
- [18] GENOVESE, C. R., ROEDER, K., AND WASSERMAN, L. (2006). False discovery control with p -value weighting. *Biometrika* **93**, 3, 509–524.
- [19] GUINDANI, M., MULLER, P., AND ZHANG, S. (2009). A Bayesian discovery procedure. *JRSS B* **71**, 905–925.
- [20] HABIGER, J. AND PEÑA, E. A. (2010). Randomized P -values and nonparametric procedures in multiple testing. *Journal of Nonparametric Statistics (iFirst article)*, 1–22. DOI: 10.1080/10485252.2010.482154.
- [21] Hoeffding, W. (1956). On the distribution of the number of successes in independent trials. *Ann. Math. Statist.* **27**, 713–721.
- [22] IHAKA, R. AND GENTLEMAN, R. (1996). R: a language for data analysis and graphics. *Journal of Computational and Graphical Statistics* **5**, 299–314.
- [23] JAMES, W. AND STEIN, C. (1961). Estimation with quadratic loss. In *Proc. 4th Berkeley Sympos. Math. Statist. and Prob., Vol. I*. Univ. California Press, Berkeley, Calif., 361–379.
- [24] JIN, J. AND CAI, T. T. (2007). Estimating the null and the proportional of nonnull effects in large-scale multiple comparisons. **102**, 478, 495–506.
- [25] KANG, G., YE, K., LIU, N., ALLISON, D., AND GAO, G. (2009). Weighted multiple hypothesis testing procedures. *Statistical Applications in Genetics and Molecular Biology* **8**, 1–21.
- [26] LAN, K. K. G. AND DEMETS, D. L. (1983). Discrete sequential boundaries for clinical trials. *Biometrika* **70**, 3, 659–663.
- [27] LANGAAS, M., LINDQVIST, B. H., AND FERKINGSTAD, E. (2005). Estimating the proportion of true null hypotheses, with application to DNA microarray data. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **67**, 4, 555–572.

- [28] LEHMANN, E. L. (1997). *Testing statistical hypotheses*, second ed. Springer Texts in Statistics. Springer-Verlag, New York.
- [29] LEHMANN, E. L., ROMANO, J. P., AND SHAFFER, J. P. (2005). On optimality of stepdown and stepup multiple test procedures. *Ann. Statist.* **33**, 3, 1084–1108.
- [30] MARSHALL, A. W. AND OLKIN, I. (1979). *Inequalities: theory of majorization and its applications*. Mathematics in Science and Engineering, Vol. **143**. Academic Press Inc. [Harcourt Brace Jovanovich Publishers], New York.
- [31] MÜLLER, P., PARMIGIANI, G., ROBERT, C., AND ROUSSEAU, J. (2004). Optimal sample size for multiple testing: the case of gene expression microarrays. *J. Amer. Statist. Assoc.* **99**, 468, 990–1001.
- [32] NEYMAN, J. AND PEARSON, E. (1933). On the problem of the most efficient tests of statistical hypotheses. *Phil. Trans. Roy. Soc. Ser. A* **231**, 289–337.
- [33] ROGOSA, D. (2003). Accuracy of api index and school base report elements: 2003 academic performance index, california department of education. *Available at* <http://www.cde.cagov/ta/ac/ap/researchreports.asp>.
- [34] ROQUAIN, E. AND VAN DE WIEL, M. A. (2009). Optimal weighting for false discovery rate control. *Electronic Journal of Statistics* **3**, 678–711.
- [35] ROYDEN, H. (1968). *Real Analysis*, Second ed. McMillan Publishing Co., Inc., New York.
- [36] RUBIN, D., DUDOIT, S., AND VAN DER LAAN, M. (2006). A method to increase the power of multiple testing procedures through sample splitting. *Stat. Appl. Genet. Mol. Biol.* **5**, Art. 19, 20 pp. (electronic).
- [37] SARKAR, S. K. (1998). Some probability inequalities for ordered MTP_2 random variables: a proof of the Simes conjecture. *Ann. Statist.* **26**, 2, 494–504.
- [38] SARKAR, S. K. (2008). Generalizing Simes’ test and Hochberg’s stepup procedure. *Ann. Statist.* **36**, 1, 337–363.
- [39] SARKAR, S. K., ZHOU, T., AND GHOSH, D. (2008). A general decision theoretic formulation of procedures controlling FDR and FNR from a Bayesian perspective. *Statist. Sinica* **18**, 3, 925–945.
- [40] SCHWARTZMAN, A., DOUGHERTY, R., AND TAYLOR, J. (2005). Cross-subject comparison of principal diffusion direction maps. *Magnetic Resonance in Medicine* **53**, 1423–1431.
- [41] SCHWEDER, T. AND SPJØTVOLL, E. (1982). Plots of p-values to evaluate many tests simultaneously. *Biometrika* **69**, 3, 493–502. <http://www.jstor.org/stable/2335984>.
- [42] SCOTT, J. AND BERGER, J. (2006). An exploration of aspects of Bayesian multiple testing. *Journal of Statistical Planning and Inference* **136**, 2144–2162.

- [43] ŠIDÁK, Z. (1967). Rectangular confidence regions for the means of multivariate normal distributions. *J. Amer. Statist. Assoc.* *62*, 626–633.
- [44] SINGH, D., FEBBO, P., ROSS, K., JACKSON, D., MANOLA, J., LADD, C., TAMAYO, P., RENSHAW, A., D’AMICO, A., RICHIE, J., LANDER, E., LODA, M., KANTOFF, P., GOLUB, T., AND SELLERS, R. (2002). Gene expression correlates of clinical prostate cancer behavior. *Cancer Cell* *1*, 302–309.
- [45] SORIĆ, B. (1989). Statistical “discoveries” and effect-size estimation. *Journal of the American Statistical Association* *84*, 608–610.
- [46] SPJØTVOLL, E. (1972). On the optimality of some multiple comparison procedures. *Ann. Math. Statist.* *43*, 398–411.
- [47] STEVENSON, R. L. (1886). *The Strange Case of Dr Jekyll and Mr Hyde*, First ed. Longmans, Green, and Co., London.
- [48] STOREY, J. (2002). A direct approach to false discovery rates. *Journal of the Royal Statistical Society, Series B* *64*, 479 – 498.
- [49] STOREY, J. (2003). The positive false discovery rate: a Bayesian interpretation and the q-value. *The Annals of Statistics* *31*, 2012 – 2035.
- [50] STOREY, J. (2007). The optimal discovery procedure: a new approach to simultaneous significance testing. *Journal of the Royal Statistical Society, Series B* *69*, 347–368.
- [51] STOREY, J., DAI, J., AND LEEK, J. (2007). The optimal discovery procedure for large-scale significance testing, with applications to comparative microarray experiments. *Biostatistics* *8*, 414–432.
- [52] STOREY, J. D., TAYLOR, J. E., AND SIEGMUND, D. (2004). Strong control, conservative point estimation and simultaneous conservative consistency of false discovery rates: a unified approach. *J. R. Stat. Soc. Ser. B Stat. Methodol.* **66**, 1, 187–205.
- [53] SUN, W. AND CAI, T. (2007). Oracle and adaptive compound decision rules for false discovery rate control. *Journal of the American Statistical Association* *102*, 901–912.
- [54] TURNBULL, B. (2006). Best proteomics data. Available at www.stanford.edu/people/brit.turnbull/BESTproteomics.pdf.
- [55] WASSERMAN, L. AND ROEDER, K. (2006). Weighted hypothesis testing. Tech. rep., Carnegie-Mellon University. (<http://arxiv.org/abs/math.ST/0604172>).
- [56] WESTFALL, P. AND TROENDLE, J. (2008). Multiple testing with minimal assumptions. *Biometrical Journal* *50*, 1–11.
- [57] WESTFALL, P. AND YOUNG, S. (1993). *Resampling-Based Multiple Testing: Examples and Methods for P-Value Adjustment*. Wiley, New York.
- [58] WESTFALL, P. H., KRISHEN, A., AND YOUNG, S. S. (1998). Using prior information to allocate significance levels for multiple endpoints. *Statistics in Medicine* *17*, 2107–2119.